

# The functional microbiome of grapevine throughout plant evolutionary history and lifetime

Paola Fournier<sup>a,b,†</sup>, Lucile Pellan<sup>b,†</sup>, Didac Barroso-Bergadà<sup>c</sup>, David A. Bohan<sup>c</sup>, Thierry Candresse<sup>d</sup>, François Delmotte<sup>b</sup>, Marie-Cécile Dufour<sup>b</sup>, Virginie Lauvergeat<sup>e</sup>, Claire Le Marrec<sup>f</sup>, Armelle Marais<sup>d</sup>, Guilherme Martins<sup>f</sup>, Isabelle Masneuf-Pomarède<sup>f</sup>, Patrice Rey<sup>g</sup>, David Sherman<sup>h</sup>, Patrice This<sup>i</sup>, Clémence Frioux<sup>h</sup>, Simon Labarthe<sup>a,h</sup>, and Corinne Vacher<sup>a,\*</sup>

<sup>a</sup>INRAE, Univ. Bordeaux, BIOGECO, Pessac, France

<sup>b</sup>INRAE, Bordeaux Sciences Agro, ISVV, SAVE, Villenave-d'Ormon, France

<sup>c</sup>INRAE, Université Bourgogne Franche-Comté, Agroécologie, Dijon, France

<sup>d</sup>Univ. Bordeaux, INRAE, UMR BFP, Villenave d'Ormon, France

<sup>e</sup>EGFV, Univ. Bordeaux, Bordeaux Sciences Agro, INRAE, ISVV, Villenave d'Ormon, France

<sup>f</sup>Univ. Bordeaux, UMR oenologie, INRAE, Bx INP, Bordeaux Sciences Agro, ISVV, Villenave d'Ormon, France

<sup>g</sup>Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, Institut des Sciences Analytiques et de Physicochimie pour l'Environnement et les Matériaux—UMR 5254, IBEAS, Pau, France

<sup>h</sup>Inria, Univ. Bordeaux, INRAE, Talence, France

<sup>i</sup>UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

\*Corresponding author: e-mail address: corinne.vacher@inrae.fr

## Contents

1. Introduction	2
2. The grapevine functional microbiome throughout evolutionary history	3
2.1 Microbiome evolution during grapevine domestication and breeding	4
2.2 Microbial interactions in the centre of origin of major grapevine pathogens	8
2.3 Microbial dispersal across geographic regions and <i>Vitis</i> species	11
3. The grapevine functional microbiome throughout plant lifetime	13
3.1 Initial microbiome and virome at the graft stage	13
3.2 Recruitment of the root microbiome from the soil reservoir	16
3.3 Seasonal assembly of the leaf microbiome in interaction with leaf pathogens	21
3.4 Dynamics of the berry microbiome to ripening and winemaking	25
3.5 Wood microbiome dysbiosis during grapevine aging and decline	28
4. Let's make the grapevine microbiome more functional	32

† Co-first authors.

4.1	Field sampling designs and statistical approaches to identify beneficial microbial taxa	32
4.2	Molecular tools to uncover the functional potential of the microbiome	35
4.3	Computational approaches to understand the grapevine holobiont as a functional and dynamic network	38
4.4	Culture-dependent approaches to validate microbial interactions and functions	44
5.	Conclusion and perspectives	45
	Acknowledgements	48
	References	48

## Abstract

European grapevine is a complex holobiont composed of two plant genomes, that of the scion (*Vitis vinifera* L.) and the rootstock (*Vitis* spp.), and a multitude of microbial genomes that collectively form the microbiome. The grapevine microbiome has been extensively described over the last decade, primarily using metabarcoding approaches. Unfortunately, metabarcoding data alone provide little information on microbial functions and outcomes of plant-microbe interactions. Here we review knowledge about the microorganisms that have a demonstrated influence, positive or negative, on the performance of the grapevine holobiont. Our review encompasses bacteria, filamentous fungi, yeasts, oomycetes and viruses. It covers aboveground and belowground microorganisms, including arbuscular mycorrhizal and ectomycorrhizal fungi. We focus on taxa and functions that protect the plant against pathogens and pests, promote growth, increase tolerance to abiotic stresses and highlight those involved in disease and decline. As the outcomes of plant-microbe interactions are labile, we examine the dynamics and functions of grapevine-microbiome interactions over both the plant lifetime and the plant evolutionary history, beginning with plant domestication. Based on the knowledge and gaps we identify, we suggest field sampling designs, culture-based experiments, molecular tools and theoretical analysis methods, including shotgun metagenomics and network models, that could be used in future research to uncover and leverage the full functional potential of the grapevine microbiome.



## 1. Introduction

European cultivated grapevine is a complex and dynamic system of plant-microbe interactions that has been shaped by humankind to produce grapes and wine. Each plant individual is typically composed of two plant genomes, that of the rootstock (*Vitis* spp.) and the scion (*Vitis vinifera* L.), onto which are linked a multitude of microbial and viral genomes. The set of microbial genomes forms the microbiome (Berg et al., 2020; Compant et al., 2019; Saikkonen et al., 2020), while the set of viral genomes forms the virome (Roossinck, 2010). The whole, including the plant

genomes, forms the hologenome (Bettenfeld et al., 2021; Theis et al., 2016; Zilber-Rosenberg and Rosenberg, 2008). This concept of each grapevine individual as being a complex and dynamic system of plant-microbe interactions (i.e., an holobiont; Bettenfeld et al., 2021; Vandenkoornhuysen et al., 2015) has not been necessary, for most of human history, to grow vines and produce wine. It could provide solutions in the future, however, to maintain grapevine cultivation in a context of climate change (Gambetta et al., 2020) and pesticide-free agriculture (Jacquet et al., 2022; Pertot et al., 2017).


The objective of this chapter is to present the state of art about the dynamics of grapevine-microbiome interactions, from the evolutionary time scale to the seasonal scale, and to propose some research avenues to increase knowledge on the functions of the grapevine microbiome. The chapter first gives an overview of the evolution of the grapevine microbiome, starting with grapevine domestication (Section 2), and then describes the microbiome dynamics during the lifetime of each plant, from the grafting stage in the nursery through to decline in the vineyard (Section 3). In both sections, we specifically highlight those microorganisms and viruses that have a demonstrated influence, positive or negative, on the performance of the grapevine holobiont (health, growth and berry quality). Based on this state of art, we suggest experimental and theoretical approaches that could lead to a better understanding of the functions of the grapevine microbiome (Section 4). We propose some experiments to identify the microbial taxa and functions that play a key role in grapevine performance under drought conditions and in agro-ecological settings. We describe how to decipher the microbial interaction networks to which these key taxa belong, to understand how they maintain in the system and regulate grapevine performance. We also provide recommendations about the biogeographic regions that could be explored in the future to isolate beneficial microbial taxa, which could then be inoculated to drive the system.



---

## 2. The grapevine functional microbiome throughout evolutionary history

This section reviews the knowledge and gaps regarding microbiome evolution during grapevine domestication and breeding, as well as the microbiome dynamics triggered by microbial dispersal across geographic regions and between members of the *Vitis* genus. Based on this state of




**Fig. 1** Area of domestication of European cultivated grapevine, *Vitis vinifera* subsp. *vinifera* (area hatched in pink), and countries where the microbiome of its wild ancestor, *Vitis vinifera* subsp. *sylvestris*, has been described (in green). Campisano et al. (2015) isolated bacterial endophytes from wild grapevine shoots (1). Ocete et al. (2015) explored taxonomic diversity of arbuscular mycorrhizal fungi in wild grapevine rhizosphere (2). Puig-Pujol et al. (2016) isolated yeast strains at the end of spontaneous fermentations of wild grapes (3). Cordero-Bueso et al. (2017) studied yeasts associated with wild grape berries (4). Asghari et al. (2019) analysed bacterial endophytes in healthy roots, stems, leaves and fruits of wild grapevines (5). Radić et al. (2021) characterised fungal communities associated with wild grapevine roots with a focus on mycorrhizae (6). For future research, we recommend exploring the centre of origin of cultivated grapevines (in pink), studying both the microbiome of wild grapevine and that of ancestral grapevine cultivars. Map created using the Free and Open Source QGIS.

art we hypothesise which geographic regions and plant genetic material are most likely to be associated with beneficial microbial taxa and functions (Figs. 1 and 2).

## 2.1 Microbiome evolution during grapevine domestication and breeding

The Transcaucasus (i.e., the geographic region between Black and Caucasian Sea, which today corresponds to Armenia, Georgia, and Azerbaijan) is the cradle of viticulture (Fig. 1). It is the most probable region



**Fig. 2** Geographic locations where the microbiome of American and Asian *Vitis* species, partially or totally resistant to pathogens of North American origin (*Plasmopara viticola*, *Erysiphe necator* and *Phyllosticta ampellicida*), has been described (green circles), as well as the native range of these resistant *Vitis* species according to Wan et al. (2013) (area hatched in pink). Brum et al. (2012) isolated foliar fungal endophytes from *V. labrusca* cv. Niagara Rosada collected in Brazil (1). Felber et al. (2016) isolated foliar endophytic fungi from grapevine cultivars Bordô and Concord (*V. labrusca*) in Brazil (2). Kernaghan et al. (2017) analysed foliar fungal endophytes of a cultivated hybrid grape variety (L'Acadie blanc) and one of its native progenitors (*V. riparia*) in eastern Canada (3). Cureau et al. (2021) explored the taxonomic diversity of berry fungal communities in *V. vinifera*, *V. rotundifolia*, *V. aestivalis*, and in hybrid *Vitis* grape varieties grown in different vineyards in Arkansas (4). Fan et al. (2020) compared endophytic fungi of *V. vinifera* cv. Red Globe (cultivated grape) with those of *V. amurensis* cv. Shuangyou (wild grape) grown in a nursery in China (5). Sun et al. (2021) studied the berry microbiome of six muscadine (*Muscadinia rotundifolia* Michx.) cultivars grown in Guangxi, China (6). Aleynova et al. (2022) compared endophytic bacterial communities of a wild grape population of *V. amurensis* in Russia with those of domesticated *V. vinifera* cultivars from Germany and California (USA) (7). Studies that focused on microorganisms isolated during or at the end of spontaneous grape fermentations are not reported on this map. For future research to identify microbial biocontrol candidates against mildews and black rot, we suggest further exploration of the microbiome of these resistant *Vitis* species in their native range (area hatched in pink), where they have co-evolved with pathogens or with related pathogens. Map created using the Free and Open Source QGIS.

of domestication of grapevine 6.000 to 8.000 years ago (This et al., 2006; Zohary and Hopf, 1993), even if secondary domestication probably occurred later in western Europe (Arroyo-García et al., 2006; Sivan et al., 2021; Terral et al., 2010). *Vitis vinifera* subsp. *vinifera*, is nowadays the main *Vitis* species used for the production of wine, table grapes and raisins worldwide. It was domesticated from a wild ancestor, *Vitis vinifera* subsp. *sylvestris* (hereafter referred to as *sylvestris*), which naturally occurs in Mediterranean Europe (Fig. 1) and southwestern Asia (Di Vecchi-Staraz et al., 2009). In the wild, *sylvestris* is a liana that climbs trees, and has very small, black and acidic berries produced by dioecious flowers. During domestication, the grape has undergone marked changes, the most important being the evolution of sex (from dioecious to hermaphroditic flowers), the increase in size of berries and bunches, and changes in berry chemical composition (This et al., 2006).

From the Transcaucasus, cultivated grape spread to Europe, first to the Mediterranean region following the main civilizations (Greeks, Romans, Etruscans, Egyptians), then to northern Europe following the arrival of Christianity. At the beginning, cultivars experienced introgressions from local wild grapes (Myles et al., 2011). Genetic flows in the other direction, from cultivated grapes to wild populations, also occurred (Di Vecchi-Staraz et al., 2009). From the medieval period onwards, cultivars evolved by crossings within *Vitis vinifera* subsp. *vinifera*, most likely in a fortuitous way at the beginning (Bowers et al., 1999), then in a more controlled manner, particularly after the emergence of mildews and phylloxera in Europe (Section 2.2). Nowadays *Vitis vinifera* genetic diversity is huge, with 6000 to 7000 cultivars in the world. However, the total number of cultivars commonly used is much lower, given that half the worldwide plantings were of just 16 varieties in 2016 (Anderson and Aryal, 2013).

In addition to genetic and phenotypic changes to the plant, grapevine domestication was accompanied by an evolution of fermentative microorganisms, particularly *Saccharomyces cerevisiae*. *S. cerevisiae* is the main agent of wine fermentation and one of the best models for understanding the eukaryotic cell (Botstein and Fink, 2011). It is naturally part of the berry microbiome (Section 3.4) and colonises many other ecological niches (Schacherer et al., 2009). *S. cerevisiae* strains naturally occurring in cultivated grapevines are often referred to as vineyard yeasts or domesticated yeasts. The comparison of vineyard and non-vineyard *S. cerevisiae* strains revealed that the oldest lineages and greatest genetic variability are found in the non-vineyard strains (Fay and Benavides, 2005). This is consistent with

the hypothesis that *S. cerevisiae* originated in natural environments and was subsequently domesticated by humans. Molecular analyses revealed that domesticated *S. cerevisiae* strains have a single origin. The initial domestication event probably occurred in Mesopotamia, concomitantly with the grapevine domestication event (Legras et al., 2007). Domestication was followed by a marked phenotypic divergence between vineyard and non-vineyard yeasts. For instance, vineyard yeasts show greater resistance to copper sulphate (Fay et al., 2004; Warringer et al., 2011), a chemical compound used as a fungicide against downy mildew since the 1880s, strongly suggesting that vineyard yeast populations have evolved in response to agricultural practices.

How has the microbiome of European cultivated grapevine, *Vitis vinifera* subsp. *vinifera*, evolved during grape domestication? Little is known for many species apart from *S. cerevisiae* (reviewed above). Several studies have suggested that grapevine domestication caused a decrease in microbiome diversity and a change in microbiome functions. Wild sylvestris grapes are inhabited by a greater diversity of endophytic bacteria, epiphytic yeasts and arbuscular mycorrhizae than domesticated grapes (Campisano et al., 2015; Cordero-Bueso et al., 2017; Ocete et al., 2015). Moreover, microbial communities of wild and domesticated grapes differ in their potential for pathogen biocontrol and plant growth promotion (Campisano et al., 2015; Cordero-Bueso et al., 2017). Finally, microorganisms generally associated with humans may have been integrated into the grapevine microbiome during domestication (Campisano et al., 2014; Yousaf et al., 2014). After the initial domestication event, the grapevine microbiome kept evolving as new varieties were created. Molecular analysis of parental relationships among grapevine varieties identified Heunisch weiss, Pinot noir N and Riesling B as the oldest and least evolved varieties, and also the main progenitors of other varieties (Lacombe et al., 2013). Microbiome taxonomic composition of Pinot and Riesling grape varieties has been compared to that of more recent varieties (Bao et al., 2022; Zhang et al., 2020). The results suggested that the compositions of rhizosphere microbial communities of Pinot and Riesling varieties were more similar to one another than those of more recent grape varieties (Bao et al., 2022). Moreover, the bacterial genera *Pseudomonas* and *Rhizobium*, which include plant growth-promoting strains (Section 3.2), were significantly enriched in the rhizosphere of Pinot noir (Bao et al., 2022). Some bacterial genera (*Bacillus*, *Turicibacter* and *Romboutsia*) were also enriched in the leaf microbiome of Pinot noir (Zhang et al., 2020).



Several studies have hypothesised that wild populations of *V. vinifera* subsp. *sylvestris* constitute a reservoir of useful microbial strains, because wild plants often harbour beneficial endophytes that are absent, or less abundant, in domesticated plants (Ofek-Lalzar et al., 2016; Sun et al., 2020). They explored the microbiome of *sylvestris* and obtained promising results. For instance, the analysis of the root microbiome showed that *sylvestris* roots were colonised by ectomycorrhizal fungi and that fungal pathogens were completely absent from roots colonised by these fungi (Radić et al., 2021). The analysis of the berry microbiome of *sylvestris* revealed a high diversity of yeasts, with some of them having promising prospects for use in oenology (Cordero-Bueso et al., 2017, 2022; Puig-Pujol et al., 2016). Yeast strains belonging to four species (*Meyerozyma guilliermondii*, *Hanseniaspora uvarum*, *H. clermontiae*, and *Pichia kluyveri*), all isolated from *sylvestris*, reduced growth of moulds caused by *Botrytis cinerea*, *Aspergillus carbonarius*, and *Penicillium expansum* (Cordero-Bueso et al., 2017). Finally, endophytic bacterial strains with biocontrol properties against several pathogens were isolated from *sylvestris* (Campisano et al., 2015), including strains belonging to *Pantoea* spp. and *Pseudomonas* spp. that showed antagonistic activity against crown gall agents (*Agrobacterium tumefaciens* and *Allorhizobium vitis*) (Asghari et al., 2019).

To identify and isolate beneficial microorganisms that may have been lost during the domestication process, future research should continue to explore the microbiome of *Vitis vinifera* subsp. *sylvestris* and that of ancestral grapevine varieties. We would recommend focusing on *sylvestris* populations in the centre of origin of cultivated grapevines (i.e., the Transcaucasus region; Fig. 1), where beneficial microbes may have co-evolved with wild progenitors, before being lost in those plants cultivated in the expanded range of grapevine. Such analysis could allow the isolation of plant growth-promoting bacteria (Gutierrez and Grillo, 2022), that might confer tolerance to abiotic stresses and grapevine pathogens of Eurasian origin (such as *Botrytis cinerea*). The case of pathogens introduced from North America is different and is discussed hereafter (Section 2.2).

## 2.2 Microbial interactions in the centre of origin of major grapevine pathogens

In the middle of the 19th century, several fungal pathogen and insect pest species of North American origin crossed the Atlantic and reached Europe, causing the destruction of a large proportion of vineyards (Gessler et al., 2011). These American species included grape phylloxera (introduced in 1863), and the agents of powdery mildew (1848), downy mildew



(1878) and black-rot (1885). Phylloxera is caused by the insect *Daktulosphaira vitifoliae*, powdery mildew by the fungi *Erysiphe necator*, downy mildew by the oomycete *Plasmopara viticola*, and black-rot by the fungi *Phyllosticta ampellicida* (asexual stage of *Guignardia bidwellii*). Scientists who established collections of wild American *Vitis* species in botanical gardens in the early 19th century probably transported these pathogen species to Europe. From there, the diseases spread to other European regions and are now reported in most wine-producing regions of the globe (Bois et al., 2017; Fontaine et al., 2021; Pirrello et al., 2019).

European viticulture was saved by the introduction of native American *Vitis* species (such as *V. riparia*, *V. rupestris* and *V. berlandieri*; Fig. 2) that were resistant to grape phylloxera. Moreover, sulphur, copper and later on, phytosanitary treatments, have been and are still used extensively to protect vines from downy and powdery mildews. The American *Vitis* species were first used to produce rootstocks resistant to phylloxera. Most European cultivated vines, planted since the end of the 19th century, are therefore grafted plants composed of a rootstock tolerant to phylloxera and a scion used to produce berries (Section 3.1). The rootstock and the scion are genetically different and may even belong to different *Vitis* species, which makes European cultivated grapevine an interesting holobiont (Biget, 2021). *Vitis* species of American and Asian origin (Fig. 2) have also been used to produce interspecific hybrids cultivated on their own roots (This et al., 2006). More recently, they have been used as sources of disease-resistance genes in breeding programmes (Merdinoglu et al., 2018) since they are not only resistant to phylloxera but also to other pests and diseases (Cadle-Davidson et al., 2011; Merdinoglu et al., 2018; Staudt and Kassemeyer, 1995; Villano and Aversano, 2020). However, these new disease-resistant varieties are already threatened by pathogen strains able to overcome resistance genes (for downy mildew: Peressotti et al., 2010, for powdery mildew: Feechan et al., 2015).

As with other plant species, the grapevine microbiome offers tremendous genetic variability that has barely been exploited in breeding programmes to date (Gopal and Gupta, 2016). What if American and Asian *Vitis* species still had something to offer to European vineyards? Although this hypothesis has yet to be tested explicitly, the higher resistance of some American *Vitis* accessions to pathogens of North American origin (Fig. 2) could be due to an increased ability of the plant to select a protective microbiome (Gopal and Gupta, 2016; Lyu et al., 2021), in addition to the acquisition of resistance genes in the plant genome. This higher disease-resistance may be the result of a longer coevolution with

the pathogens (Jürges et al., 2009). The resistance of some Asian *Vitis* species (Fig. 2) to mildews is more difficult to explain. The resistance of *V. amurensis* to downy mildew, *Plasmopara viticola*, might be due to the presence in Asia of a related pathogen species, *P. amurensis*, which has similar infection mechanisms (Jürges et al., 2009). Recent genetic analyses reopened the question of a possible origin of powdery mildew from Asia (Gur et al., 2021), and might account for the resistance of Asian *Vitis* species to powdery mildew.

Several studies have now characterised the microbiome of American and Asian *Vitis* species. They have found a high richness of endophytes (Fan et al., 2020; Kernaghan et al., 2017) and isolated strains with biocontrol properties against pathogens of North-American origin, as well as other pathogens. For instance, *V. amurensis* cv. Shuangyou in China hosted *Alternaria* strains effective against downy mildew (Musetti et al., 2006). Fungal endophytes isolated from leaves of *V. riparia* in eastern Canada showed high levels of inhibition of grey mould (caused by *Botrytis cinerea*) and black-foot disease (*Cylindrocarpon destructans*). Fungal strains inhibiting grey mould belonged to *Ramularia pratensis*, *Phoma aliena* and *Fusarium acuminatum*, while strains inhibiting black-foot belonged to *Ramularia* spp., *Phoma* spp., and *Biscogniauxia mediterranea* (Kernaghan et al., 2017). Similarly, endophytes isolated from *V. labrusca* leaves in Brazil showed in vitro antagonistic activity against several grapevine pathogens (Brum et al., 2012; Felber et al., 2016). Isolates of *Flavodon flavus* and *Colletotrichum gloeosporioides* strongly inhibited the root pathogen *Fusarium oxysporum* f. sp. *herbemontis* (Brum et al., 2012). Isolates belonging to *Phoma* spp. inhibited the growth of the causal agent of leaf blight (*Alternaria* spp.), isolates of *Fusarium culmorum* regulated anthracnose (*Sphaceloma* spp.) and *Sordariomycetes* isolates controlled ripe rot of grapes (*Glomerella* spp.) (Felber et al., 2016). The fungi *Acaromyces ingoldii*, previously reported for its plant protection functions, was found to be the most abundant taxon in the berry microbiome of muscadine grape (*Muscadinia rotundifolia* Michx.) in China (Sun et al., 2021). In addition, the bacterial genus *Rahnella* was detected in the microbiome of *M. rotundifolia* and might play a role in the biological control of grapevine crown gall, caused by *Agrobacterium vitis* (Chen et al., 2007).

To move towards pesticide-free viticulture, research on the microbial biocontrol of diseases of North American origin (powdery mildew, downy mildew and black-rot) should be fostered. To identify candidate biocontrol agents, we would recommend analysing the microbiome of American *Vitis* species in their native range (Fig. 2), where they have co-evolved with the

pathogens and the rest of the microbiome. It could also be relevant to analyse the microbiome of resistant Asian *Vitis* species (Fig. 2) in areas where they have co-evolved with pathogens related to pathogens of North-American origin. The level of disease-resistance should first be evaluated and the microbiome of all plant compartments colonised by pathogens should be characterised. To highlight microbial taxa that play a role in disease resistance (Section 4.1), the microbiomes of resistant accessions could then be compared to that of sensitive ones in the same geographic region. The inference of microbial interaction networks in resistant *Vitis* species could also generate hypotheses about the microbial taxa or consortia conferring disease resistance (Section 4.3).

### 2.3 Microbial dispersal across geographic regions and *Vitis* species

As described above, cultivated grapevine is a complex holobiont that incorporates plant and microbial genomes of European, American and Asian origin (Sections 2.1 and 2.2). To understand how this complex system has assembled and evolved since grapevine domestication (Section 2.1), it is necessary to understand the origin and migratory routes of grapevine-associated microbes. To date, research has focused on the migratory flows of the oenological yeast *Saccharomyces cerevisiae* as well as those of major pathogens of grapevine (*Plasmopara viticola*, in particular).

Molecular evidence for the historical presence of *Saccharomyces cerevisiae* in wine fermentation was obtained from pottery jars hidden in the tomb of King Scorpio I, one of the earliest kings of Egypt, dating back to 3150 BC (Cavaleri et al., 2003). At that time, fermentations were only spontaneous (Marsit and Dequin, 2015), suggesting that *S. cerevisiae* was already an integral part of the berry microbiome. Legras et al. (2007) analysed migration routes of *S. cerevisiae* and showed that they correspond to the known migration routes of cultivated grapevine (Grassi and De Lorenzis, 2021). From Lebanon, yeast migration may have taken place across the Mediterranean region towards Italy, Spain and France, through three main routes: (1) the Rhône Valley-Burgundy-Alsace and Nantes route that goes along the Rhône and Loire rivers; (2) the Italy-Cognac route; and, (3) the Danube Valley route. This migration of vineyard yeast strains is accounted for, at least in part, by the trade of grape varieties. For instance, the Muscadet grape variety was imported from Burgundy to Nantes during the 15th century, while Ugni blanc, the dominant grape variety in the

French Cognac region, originated in Italy. Genetically-distant Austrian yeast strains may have been domesticated from local wild *sylvestris* populations (Section 2.1).

*Plasmopara viticola* is a complex of five species, each with a unique degree of host specialisation within the American Vitaceae (Fontaine et al., 2021; Rouxel et al., 2013, 2014). Fontaine et al. (2013, 2021) found that all invasive grapevine downy mildew populations worldwide belonged to the small clade *P. viticola* f. sp. *aestivalis*, which parasitizes *Vitis aestivalis* in Northeast America (Fig. 1). *P. viticola* was first introduced into Western Europe in the middle of the 19th century (Section 2.2), and spread to Central and Eastern Europe. Fontaine et al. (2013) and Gobbin et al. (2006) revealed, using population genetic methods, a low genetic diversity and weak population structure for *P. viticola* across European vineyards, suggesting that the strains initially introduced into Europe came from a single source population of North American origin. European populations of downy mildew then served as a source for secondary introductions into other viticultural regions of the world, including China, South Africa, and Australia. A third introduction event occurred later, that spread the disease from Australia to Argentina.

The migration routes of *Erysiphe necator* and *Phyllosticta ampellicida* have been less studied than those of *Plasmopara viticola*. According to Brewer and Milgroom (2010), populations of *E. necator* in Europe are derived from two separate introductions from the Eastern United States. These initial introductions into Europe were followed by secondary introductions from Europe into the Western United States and Australia. Recently, Gur et al. (2021) proposed an additional non-American (possibly Asian) origin of *E. necator*. There is even less knowledge of the invasion pathways of *P. ampellicida* (Narduzzi-Wicht et al., 2014). Rinaldi et al. (2017) suggested that the pathogen was introduced into Northeastern Italy from Eastern Europe and that two introductions occurred in Portugal, one from France or Italy, and another from America. Interestingly, these American pathogens not only colonised the European cultivated grapevine *Vitis vinifera* subsp. *vinifera*, but were also incorporated into the microbiome of its wild ancestor *Vitis vinifera* subsp. *sylvestris* (Section 2.1). As a result, the distribution of wild grapevines has been greatly reduced over the last 150 years and it is now considered an endangered plant species in Europe (Arnold et al., 2005).

Future research should continue unravelling the migratory routes of grapevine pathogens and extend these efforts to other member of the

functional microbiome (Section 3). This is crucial to identify those geographic locations where grapevine–pathogen–microbiome co-evolution has occurred, as well as the *Vitis* species involved in this coevolutionary process (Fig. 2). This knowledge is necessary to design microbiome studies informed by the biogeography and evolutionary history of grapevine. Microbiome dynamics on a smaller temporal scale should also be considered and is discussed below (Section 3).




### 3. The grapevine functional microbiome throughout plant lifetime

This section describes how the microbiome assembles, functions and changes over the course of grapevine life. It highlights the plant–microbe interactions that have a proven, direct and significant influence on the grapevine performance (growth, health and berries quality) at every life stage (Fig. 3). The section deals with both the microbiome (bacteria, filamentous fungi, yeasts and oomycetes) and the virome, including phytoviruses (plant-infecting viruses), mycoviruses (fungi-infecting viruses) and phages (bacteria-infecting viruses). It covers both aboveground and belowground microorganisms, including arbuscular mycorrhizal and ectomycorrhizal fungi.

#### 3.1 Initial microbiome and virome at the graft stage

The life of cultivated grapevine generally begins in a nursery with a graft (Fig. 3). The two components of the graft, the scion and the rootstock, are produced in nursery “mother” plots. Mother plants produce stems that are pruned during the dormant stage in winter to make cuttings that are then stored in a cold room. The rootstock and scion cuttings and the grafted plants then undergo different treatments depending on the country and the nursery (Gramaje and Armengol, 2011). Most often, immediately after grafting, the scion and the graft zone are dipped in a wax solution, which may contain plant growth regulators (mainly auxins) and fungicides. The grafted plants are then incubated for 2–3 weeks under controlled temperature (28°C) and high humidity (90%), in order to promote callusing. A second waxing step is then performed, before planting in pots in the greenhouse, or in nursery fields. Potted plants can be shipped directly to winegrowers and planted in the field next autumn, while plants kept in nursery fields are usually shipped as 1-year-old, bare-root plants to be planted in spring.



**Fig. 3** Microbial functional groups that colonise cultivated grapevine from the nursery to plant decline, as well as ecological interactions (black arrows) and microbial fluxes (yellow arrows) that drive the microbiome dynamics during plant lifetime. Plant-microbe-microbe ecological interactions form complex networks (Section 4) that could not be represented here for clarity. Only a few ecological interactions are represented. For the same reason, we could not represent all the fungal (F), bacterial (B), oomycete (O) species and viruses (V) that significantly influence grapevine performance and wine-making process. We only represented a few well-studied species or species groups. AAB, Acetic Acid Bacteria; GFLV: Grapevine Fanleaf Virus; GLRV, Grapevine Leafroll associated Virus; Other species names: *Agrococcus baldri*, *Allorhizobium vitis*, *Aureobasidium pullulans*, *Bacillus subtilis*, *Botrytis cinerea*, *Burkholderia phytofirmans*, *Candidatus phytoplasma*, *Erysiphe necator*, *Eutypa lata*, *Ilyonectria liriodendri*, *Neofusicoccum parvum*, *Oenococcus oeni*, *Pantoea agglomerans*, *Phyllosticta ampellicida*, *Plasmopara viticola*, *Pseudomonas fluorescens*, *Pythium oligandrum*, *Rhizopagus irregularis*, *Saccharomyces cerevisiae*, *Xylella fastidiosa*.

The many steps of the grafting process are critical for assuring the quality of the plants produced. First, the choice of the two plant genomes is crucial. Not accounting losses due to incompatibility between genotypes (Tedesco et al., 2022), about 50% of the plants produced are unmarketable because of the low quality of the graft (Carrere et al., 2022). Second, it is fundamental for nurseries to produce plants that do not carry diseases (Waite et al., 2015), in particular phytoviruses. Viruses trigger both graft incompatibility and decline of young vines. For instance, the transmission of grapevine leafroll-associated virus 2 (GLRaV-2; Fig. 3) from scion to rootstock results in graft incompatibility (Rowhani et al., 2017). Rootstock susceptibility to infection depends on the genotype, with genotypes 1616 Couderc, Kober 5BB, 1103 Paulsen being the most susceptible rootstocks (Alkowni et al., 2011; Rowhani et al., 2017; Uyemoto et al., 2000, 2001). Reverse incompatibility, where the rootstock is the source of a latent virus, has not been observed in grapevine yet. The decline of young vines triggered by joint infection with GLRaV-2 and grapevine virus B (GVB) infection, or GLRaV-1 and grapevine virus A (GVA), was also observed (Rowhani et al., 2017). The mechanisms underlying such declines have not been clearly determined yet, but they might involve the production of viral small interfering RNAs (vsiRNAs) by the plant cell machinery, which in turn would affect plant gene expression (Smith et al., 2011). Management of these phytoviruses relies primarily on preventive measures (such as the use of clean propagation and planting materials) and robust diagnostics (Maliogka et al., 2015; Martelli, 1999). In addition, hot water treatment (HWT) of the grafted plant (Eichmeier et al., 2018) is sometimes performed to reduce infection by phytoplasmas (flavescence dorée, bois noir, grapevine yellows) and by fungi responsible for grapevine trunk diseases (Section 3.5). *Trichoderma* spp., which are soil microbes with an antagonistic activity against some pathogenic fungi, are sometimes inoculated by nurseries (Eichmeier et al., 2018; Jaarsveld et al., 2021). Finally, mycorrhiza (*Rhizophagus irregularis*, in particular) can also be inoculated at the nursery stage (Fig. 3), to increase grapevine resistance to biotic and abiotic stresses by promoting the structure and development of the root system (Bettenfeld et al., 2021).

All these manipulations and treatments of scions and rootstocks in the nursery can influence the initial grapevine microbiome and its future trajectory of assembly. Gramaje et al. (2022) analysed the fungal communities associated with rootstocks harvested in two mother plots located 800m apart. No difference in fungal community composition between these



two locations was detected, but the fungal communities varied along the grafting process (after cutting from mother plants, after cold storage and hydration, after grafting and callusing, and in dormant grafted plants). The core mycobiome was composed of the genera *Cadophora*, *Cladosporium*, *Penicillium* and *Alternaria* in both rootstocks. Numerous genera associated with grapevine trunk diseases (*Cadophora*, *Dactylonectria*, *Diaporthe*, *Diplodia*, *Ilyonectria*, *Neofusicoccum*, *Phaeoacremonium* and *Phaeomoniella*; Section 3.5) and potential biocontrol activities (*Aureobasidium pullulans*) were identified (Fig. 3). The pathogenic genus *Neofusicoccum* was persistent throughout the propagation process (Fig. 3), confirming the role of rootstock as a primary source of many pathogens. Lade et al. (2022) compared the contribution of nursery, scion variety and rootstock variety to fungal community composition in the graft zone and found that the nurseries (i.e., plant production practices) played a major role, followed by rootstocks and scion varieties.

To better understand and manage the assembly trajectory of the grapevine microbiome, special attention should be paid to the production practices of grafted plants in nurseries. We recommend using the framework of microbial community coalescence (Rillig et al., 2015) to analyse the influence of grafting on the microbiome. The microbiome of the grafted plant would then be considered as a joining of previously separate communities, forming a new entity that is not easily separable and with properties distinct from the progenitor communities (Rillig et al., 2015). At the graft stage, major physiological changes occur (Cookson et al., 2013; Prodhomme et al., 2019), including vascular connection between rootstock and scion that influences sap flow and associated fluxes of microorganisms (Deyett and Rolshausen, 2020). These physiological processes have barely been studied although they could have a significant impact on the initial microbiome of grafted plants and, through priority effects (Zhou and Ning, 2017), on the recruitment of the microbiome after planting (Section 3.2).

### 3.2 Recruitment of the root microbiome from the soil reservoir

Winegrowers usually receive grafted vines (potted or bare-rooted) from nurseries (Section 3.1) for planting in their vineyards. The young plants then begin to recruit microorganisms from the soil (Fig. 3), which is the major reservoir of microorganisms for the plant microbiome (Griggs et al., 2021; Liu et al., 2019; Swift et al., 2021). Recruitment is based on the active release by the roots of a wide range of carbon-containing compounds known as rhizodeposits (Chaparro et al., 2014; del Martínez-Diz et al., 2019). The rhizodeposits represent nearly 40% of the photosynthates

produced by a plant (Bais et al., 2006). These compounds serve as a substrate for the development of microorganisms and make the rhizosphere (i.e., the transition zone between soil and plant roots) much more attractive for microbes than surrounding soil. The nature and quantity of rhizodeposits depend on the plant genotype and vary across seasons and throughout plant life, allowing the plant to actively shape the composition of rhizosphere microbial communities (Berlanas et al., 2019). Beneficial microorganisms may for instance be recruited under stressful conditions, a mechanism known as the “cry for help” reaction (Rizaludin et al., 2021). Some rhizospheric microorganisms develop physical interactions with the root (case of mycorrhizae, for instance) and some of them can even enter the root endosphere. Once in the root endosphere, some microorganisms colonise the vascular tissues and disseminate in the aerial parts of the plant (Fig. 3), sometimes reaching leaves and berries (Darriaut et al., 2022; Liu et al., 2019). Not all microorganisms can colonise the plant internal tissues from surrounding soil, which creates a decreasing gradient of microbial diversity from the outside of the root towards the inside (del Martínez-Diz et al., 2019; Marasco et al., 2018; Zarraonaindia et al., 2015).

Many studies have examined the taxonomic composition of the grapevine rhizosphere and root microbiome. The same major taxonomic groups are generally represented. Pseudomonadota, Actinomycetota and Acidobacteriota (formerly named Proteobacteria, Actinobacteria and Acidobacteria; Oren and Garrity, 2021) are the dominant bacterial phyla, followed by Bacillota and Bacteroidota (i.e., Firmicutes and Bacteroidetes) (Bettenfeld et al., 2021; Dries et al., 2021; Novello et al., 2017; Zarraonaindia et al., 2015). Ascomycota, followed by Basidiomycota, are the dominant fungal phyla (Bettenfeld et al., 2021; Liu and Howell, 2021). At a finer taxonomic scale, the microbiome composition varies with soil physical and chemical features (Bettenfeld et al., 2021; Signorini et al., 2021), geography and climate (Hernandez and Menéndez, 2019; Liu et al., 2019), plant age and rootstock genotype (Berlanas et al., 2019; Ji et al., 2019), and vineyard management practices (Likar et al., 2017) including applications of plant protection products and fertilizers (Canfora et al., 2018; Zaller et al., 2018) and the presence of vegetation cover (Vukicevich et al., 2018).

Root-microbe interactions can foster vine growth and productivity through various mechanisms. Plant Growth-Promoting (PGP) microbes (Fig. 3) can solubilise and make available nutrients to the roots (Darriaut et al., 2022). The mineral elements needed by the plant (phosphorus, nitrogen and phosphate) are naturally present in the soil but cannot be efficiently

or directly assimilated by plants because they are complexed with other molecules. Other PGP bacteria produce enzymes (phosphatases) or organic acids (e.g., hydrogen cyanide; [Rijavec and Lapanje, 2016](#)) that solubilise inorganic phosphate, while others produce ammonium ( $\text{NH}_4^+$ ) or convert nitrites ( $\text{NO}_2^-$ ) to nitrates ( $\text{NO}_3^-$ ), that are easily absorbed by the plant. Bacteria, including *Bacillus herbersteinensis*, *Bacillus licheniformis*, *Micrococcus* spp., *Pseudomonas* spp., and *Pantoea agglomerans*, can combine these mechanisms ([Asghari et al., 2020](#); [Baldan et al., 2015](#)). In the grapevine rhizosphere, the bacterial genera *Pseudomonas*, *Streptomyces* and *Rhizobia* are those most involved in phosphorus and nitrogen cycles ([Bona et al., 2018](#)). Some fungal genera are also involved in phosphate solubilisation, such as the genus *Mortierella* ([Carbone et al., 2021](#); [Liu and Howell, 2021](#)). Rhizosphere PGP bacteria ([Fig. 3](#)) can also produce hormones, such as auxin, which directly stimulates plant growth ([Morales-Cedeño et al., 2021](#)). For instance, *Agrococcus baldri* and *Paenibacillus* spp. produce hormones that, even at low concentration, cause root elongation, branching and increase root diameter and density ([Baldan et al., 2015](#)). Bacterial species, such as *Bacillus amyloliquefaciens*, can produce microbial siderophores, which bind with iron, and contribute to plant iron nutrition ([Mardanov et al., 2019](#)). Other endophytic bacteria associated with root tissues (*Pseudomonas* spp., in particular), produce enzymes (ACC-deaminases) that cleave ACC (1-aminocyclopropane-1-carboxylic acid), which is the immediate precursor of ethylene in plants. This reduces the production of ethylene and limits the damage caused by biotic and abiotic stresses ([Marasco et al., 2018](#); [Saraf et al., 2010](#)).

Arbuscular mycorrhizal fungi (AMF) also play an important role in water and nutrient absorption. Belonging mainly to the *Glomeromycota* phylum, these endomycorrhizal fungi establish symbioses with vine roots ([Balestrini et al., 2010](#)). Hyphae increase the exchange surface between the vine roots and the soil. They also activate phosphorus and nitrogen transporters in root cortical cells ([Trouvelot et al., 2015](#)). AMF are the most abundant mycorrhizae associated with grapevine roots and include species belonging to the genera *Acaulospora*, *Diversispora*, *Funneliformis*, *Glomus* and *Rhizophagus* ([Darriaut et al., 2022](#); [Liu and Howell, 2021](#)). They are actively recruited by vine roots during water stress ([Carbone et al., 2021](#)).

Grapevine roots are not only colonised by beneficial microorganisms such as PGP and AMF, but also by bacterial, fungal and viral pathogens ([Fig. 3](#)). The bacterial pathogen *Allorhizobium vitis*, the causal agent of crown

gall, is frequently isolated from diseased plants in the vineyard (Habbadi et al., 2020). The bacteria harbours tumour-inducing plasmids (pTi), part of which integrate into the plant genome (Transfert-DNA, T-DNA) to induce the development of root or crown galls and the synthesis of opines, which the bacteria utilises as nutrients. The spread of pTI plasmids can be facilitated by bacteria belonging to the *Novosphingobium* genus, which are therefore pathogen stimulants (Gan et al., 2019). Black-foot disease, caused by *Ilyonectria* spp. (*I. liriodendri*, in particular), is responsible for root necrotic lesions, with a reduction in root biomass, besides other symptoms that include reduction of internodes, loss of vigour, patchy foliage and smaller leaves (Bleach et al., 2021). The fungal genus *Ilyonectria* is found in many metabarcoding analyses of vineyard soil because it can persist for several years after the removal of infected vines (de Brito et al., 2019; Deyett and Rolshausen, 2020; Liu and Howell, 2021; Rivas et al., 2022). Several fungal genera associated with grapevine trunk diseases (*Botryosphaeria* and *Diplodia*; Section 3.5), Petri disease (*Cadophora*, *Phaeoacremonium* and *Phaeoconiella*) and Diaporthe dieback (*Diaporthe*) have been detected in asymptomatic root tissues (del Martínez-Diz et al., 2019; Deyett and Rolshausen, 2020; Liu and Howell, 2021; Rivas et al., 2022). *Candidatus phytoplasma* spp., the causal agent of bois noir disease, was also detected in healthy root tissues (Marasco et al., 2018), indicating that the root endosphere may be an important reservoir for latent pathogens (del Martínez-Diz et al., 2019; Liu and Howell, 2021). Finally, grapevine roots can be infected by phytoviruses, some of which are transmitted by soil nematodes. For instance, grapevine fanleaf virus (GFLV) and arabis mosaic virus (ArMV), the main causal agents of grapevine fanleaf disease, are transmitted by the nematode species *Xiphinema index* and *Xiphinema italiae* in the case of GFLV, and *Xiphinema diversicaudatum* in the case of ArMV (Oliver and Fuchs, 2011). The nematodes feed on the tips of growing roots by pricking them with a stylet. By successively feeding on neighbouring vines, they disseminate the virus retained in their oesophageal tracts (Maliogka et al., 2015), producing the often patchy distribution of virus-infected plants.

The root microbiome also harbours microorganisms that directly inhibit pathogen growth, through antagonistic microbial interactions, or indirectly limit disease severity or frequency by stimulating the plant immune system or by acting on vector populations (Fig. 3). Microbial interactions responsible for pathogen biocontrol include antibiosis, hyperparasitism, spatial and nutritional competition, interference with pathogen signalling, and disease

symptom reduction by degradation of pathogen toxins or virulence factors (Compant et al., 2013). For instance, the biocontrol activity of *Bacillus amyloliquefasciens* mainly relies on antibiosis. This bacteria synthesises a wide range of antimicrobial compounds (surfactin, plantazolicin, macrolactin, bacillaene, fengycin, diffidin, bacillibactin, and bacilysin) that regulate the growth of several grapevine pathogen species, including *Botrytis cinerea*, *Phaeoacremonium aleophilum*, *Phaeoconiella chlamydospora* (Mardanov et al., 2019). Root microbes also stimulate plant defences, by inducing oxidative bursts, activating signalling pathways (salicylic acid, ethylene/jasmonic acid), and increasing the synthesis of pathogenesis-related (PR) proteins, lytic enzymes, polyamines and phytoalexins (Héloir et al., 2019). This stimulation increases the capacity of plants to react to ongoing infections, or to future ones by priming the immune system. Bacteria of the *Burkholderia* genus increase the production of PR proteins when inoculated on the roots of grapevine infected with *B. cinerea* (Esmaeel et al., 2020), and the strain *Burkholderia phytofirmans* PsJN can migrate to the aerial parts of grapevine and form biofilms that directly limit the spread of *B. cinerea* (Miotto-Vilanova et al., 2016). Many other root microorganisms have a proven antagonistic effect on grapevine pathogens but their mode of action is not always known. Other bacteria with a biocontrol activity include *Streptomyces* spp., that reduce young grapevine decline and black-foot disease (González-García et al., 2019; Pilar Martínez-Diz et al., 2021), and *Bacillus subtilis*, which directly limits the growth of fungal pathogens related with grapevine trunk diseases (Berlanas et al., 2019). The latter also stimulates plant defences, similar to *Pseudomonas fluorescens*, *Pantoea agglomerans*, *Acinetobacter lwoffii* (Aziz et al., 2016, 2020; Magnin-Robert et al., 2007). Root fungi, in particular AMF, also play a role in disease biocontrol. AMF of the *Glomus* genus and *Rhizophagus irregularis* reduce the severity of black-foot disease (Berlanas et al., 2019; Darriaut et al., 2022). AMF may also protect grapevines from GFLV by inhibiting the proliferation and penetration of its nematode vector (*X. index*) in roots (Hao et al., 2018). In addition, the fungal species *Trichoderma atroviride* and *Aureobasidium pullulans* are antagonists of the fungal pathogens associated with Petri disease (*Phaeoconiella chlamydospora* and *Phaeoacremonium minimum*), and of *Diplodia seriata* which is associated with Botryosphaeria dieback (del Martínez-Diz et al., 2019). *T. atroviride* also fosters plant defences (Stempien et al., 2020). Finally, the oomycete *Pythium oligandrum* has mostly indirect negative effects on fungal pathogens associated with Petri disease (Yacoub et al., 2016).

This state of art demonstrates that grapevine roots constitute a rich microbial compartment that is quite well understood from a functional point of view. Many microorganisms that have a positive (such as PGP bacteria, AMF and biocontrol agents) or negative (such as *Allorhizobium vitis* or phytoviruses) influence on grapevine have been identified and their mode of action has been described. Shotgun metagenomics and metabolomics associated with metabolic network modelling (Section 4.3) could further advance this understanding by allowing us to build predictive models of the nutritional interactions between grapevine and microbes in the rhizosphere (Jacoby et al., 2017). These models could inform the design of microbial consortia to inoculate to boost plant growth and defence. To optimise soil management, future research should focus on belowground-aboveground microbial interactions in vineyards. They could first assess the effects of soil management on soil microbiome (Bettenfeld et al., 2021; Signorini et al., 2021), and then the cascading effects of the soil microbiome on the root, trunk, leaf and berry microbiomes (Sections 3.3 and 3.4) and their consequences on grapevine health and wine quality.

### 3.3 Seasonal assembly of the leaf microbiome in interaction with leaf pathogens

The grapevine leaf microbiome is dynamic: it assembles in spring, changes during the vegetative season and then disassembles during leaf senescence. In spring, the buds burst and let the first shoots appear. As vines are lianas, the shoots hang on to the trellis with their tendrils and grow by producing new leaves, at a rate of about one every 3 days in the Bordeaux area, France (Calonnec et al., 2008). At a given time, each vine thus displays a mosaic of leaves of different ages. The leaves at the distal end of shoots are the youngest and the leaves near the stock are the oldest (see Fan et al., 2020). As the shoot grows, the young leaves expand, become mature, and are colonised by epiphytic microorganisms (the phyllosphere microbiome *sensu stricto*; Behrens and Fischer, 2022; Vacher et al., 2016a) and endophytic microorganisms (Pacífico et al., 2019) that collectively form the leaf microbiome. The latter is largely dominated by fungi belonging to the Ascomycota and by bacteria belonging to the Actinomycetota, followed by Pseudomonadota and Bacillota (Bettenfeld et al., 2021; Fort et al., 2016; Oren and Garrity, 2021; Singh et al., 2018). The fungal species *Aureobasidium pullulans* was dominant at the end of growing season in several independent metabarcoding studies, representing more than half of the fungal community in old leaves (Grube et al., 2011; Knapp et al., 2021).

Leaf microbiome dynamics depends on many factors, and first and foremost on the specific growth pattern of grapevine (Calonnec et al., 2008, 2018). Indeed, the date at which a leaf emerges determines its morphology and physiology (Chitwood et al., 2016; Majer and Hideg, 2012) and also the microbial fluxes and environmental perturbations to which it will subsequently be exposed. Leaves that emerge first are colonised by microorganisms that overwintered in buds or on woody parts (case of primary infections by the fungal pathogen *Erysiphe necator*, Gadoury et al., 2012), or in senescent leaf debris that remained on the soil (case of primary infections by the oomycete pathogen *Plasmopara viticola*, Burruano, 2000). They are also colonised by microorganisms circulating in the xylem or in the phloem (Fig. 3). These circulating microorganisms may be beneficial, as in the case of the bacterium *Burkholderia phytofirmans* (Compant et al., 2008) or they may be pathogens, like the bacterial pathogen *Xylella fastidiosa* (Deyett and Rolshausen, 2019) and viruses and phytoplasmas (Laimer et al., 2009; Zherdev et al., 2018). Wind and rain disseminate microorganisms from multiple sources (vines, soil, adjacent semi-natural and natural habitats) (Fig. 3) and may also contribute to the initial colonisation of the phyllosphere (Abdelfattah et al., 2019; Griggs et al., 2021). By contrast, those leaves that emerge later in the season are in direct contact with other grapevine leaves that already have a more developed microbiome. Leaf-to-leaf transmission of the microbiome (Fig. 3), which may transiently involve the atmosphere or other plant organs, is the dominant microbial colonisation pathway in grapevine leaves (Abdelfattah et al., 2019). This transmission is responsible for secondary infection cycles in leaf pathogens. Late-emerging leaves also face greater activity from fauna (arthropods, birds) that are underestimated microbial vectors (Griggs et al., 2021; Stefanini and Cavalieri, 2018). Finally, it should be noted that leaves of different ages do not provide the same habitat for microorganisms. Leaf age determines the structure of the cuticle and epicuticular waxes, the amount of available nutrients, the concentration of secondary metabolites, all of which regulate the growth of microbial populations (Griggs et al., 2021; Shakir et al., 2021). Leaf age also determines immune status as microbial colonisation can stimulate defences and prime the leaf to be more resistant to subsequent infection (Chaudhry et al., 2021; Shakir et al., 2021; Section 3.2). The effect of leaf age on disease development is well understood and is termed ontogenic resistance. It has been demonstrated for several grapevine leaf pathogens (Calonnec et al., 2018; Qiu et al., 2015; Steimetz et al., 2012). All of these factors lead to



intra-canopy variation in the composition of the grapevine leaf microbiome (Calonnec et al., 2008; Fan et al., 2020) as well as strong seasonal dynamics. Seasonal dynamics, however, are less pronounced in bacterial communities than in fungal communities, whose diversity decreases markedly over the growing season (Fort et al., 2016; Gobbi et al., 2020; Liu and Howell, 2021; Pinto et al., 2014; Singh et al., 2018).

In addition to seasonal intra-canopy dynamics, the grapevine leaf microbiome exhibits high spatial variability. Its composition changes significantly from one geographical region to another (Oliveira et al., 2018; Singh et al., 2018) and from one vineyard plot to another (Barroso-Bergadà et al., 2021, but see Knapp et al., 2021). These spatial variations are explained by vine-related (scion variety, cultural practices) and environmental factors (climate and microclimate, landscape structure) (Bettenfeld et al., 2021). Several studies have attempted to establish a hierarchy between these factors. For example, variety has a significant effect on leaf microbiome composition (Singh et al., 2018, 2019; Zhang et al., 2020), but its effect is much smaller than that of geographic and climatic factors (Singh et al., 2018). The effect of phytosanitary treatments appears to be less important than seasonal dynamics (Gobbi et al., 2020), in agreement with results demonstrating high resilience of the phyllosphere microbiome to phytosanitary treatments (Perazzolli et al., 2014). The cropping system as a whole, however, has a substantial influence on the diversity and composition of leaf microbiome, with significant differences found between organic and conventional farming (Barroso-Bergadà et al., 2021; Castañeda et al., 2018; Kernaghan et al., 2017; Miura et al., 2019). Finally, several studies investigated whether forest patches or orchards adjacent to grapevine plots structure the grapevine leaf microbiome, with varying results (Castañeda et al., 2018; Castañeda and Barbosa, 2017; Chandra et al., 2020; Fort et al., 2016; Miura et al., 2019).

Among the multitude of microorganisms that colonise grapevine leaves, some affect plant health and growth. Pathogens and their mode of action have been the most studied (Armijo et al., 2016). Powdery and downy mildews (respectively caused by the fungal species *Erysiphe necator* and the oomycete species *Plasmopara viticola*; Fig. 3) are the most important diseases of grapevine (Bois et al., 2017). Black-rot (caused by the fungi *Phyllosticta ampellicida*, Fig. 3) is a foliar disease that is becoming increasingly important in Europe, concomitant with efforts to reduce fungicide use (Molitor and Beyer, 2014). All three pathogens are introduced species of

North-American origin (Sections 2.2 and 2.3). When these pathogen species colonise the grapevine leaf, they encounter the resident microbiome and virome, which can act as a barrier to their growth through direct ecological interactions (Chaudhry et al., 2021; Hacquard et al., 2017), such as exploitative competition, interference competition (antibiosis), and hyperparasitism. Once this first barrier is overcome, pathogens are confronted by the plant's immune system, which is itself stimulated by the microbiome, especially the root microbiome (Section 3.2).

Natural antagonists of downy mildew (Fig. 3) have been identified using culture-dependent methods. Several bacterial and fungal species, isolated from grapevine leaves or previously detected in the leaf microbiome, have a negative effect on *Plasmopara viticola* under experimental conditions. For fungi these include *Acremonium byssoides*, *Alternaria alternata*, *Beauveria bassiana*, *Epicoccum nigrum* (Burruano et al., 2008, 2016; Kortekamp, 1997; Musetti et al., 2006; Rondot and Reineke, 2019; Taguiau et al., 2021; Zanzotto and Morroni, 2016), and several *Fusarium* species (Bakshi et al., 2001; Falk, 1996). The genus *Bacillus*, including *B. subtilis* and *B. pumilus*, predominates among *P. viticola* antagonistic bacteria (Bruissson et al., 2019; Dagostin et al., 2011; Zanzotto and Morroni, 2016; Zhang et al., 2017). Moreover, root colonisation by the arbuscular mycorrhizal fungus *Rhizophagus irregularis* (Sections 3.1 and 3.2) increases grapevine resistance to *P. viticola* in leaves (Cruz-Silva et al., 2021). Finally, the virome of *P. viticola* has been analysed using metatranscriptomics in search for hypovirulence-inducing mycoviruses (Chiapello et al., 2020). However, despite the description of many new mycoviruses (283 new viruses associated with lesions caused by *P. viticola*), the limited availability of phenotypic data on fungal hosts has so far prevented the association of particular viruses with variations in fungal virulence.

Pathogen-microbiome-virome interactions are not well documented for powdery mildew (Panstruga and Kuhn, 2019), but some antagonistic interactions (Fig. 3) have been demonstrated experimentally. Several natural hyperparasites of *Erysiphe necator* have been identified among the *Ampelomyces*, *Lecanicillium*, and *Acremonium* genera (Falk, 1995; Ghule et al., 2019; Kiss, 2003). As with downy mildew, powdery mildew growth is reduced by *B. subtilis* and *B. pumilus* (Compant et al., 2013; Crisp et al., 2006). Network inference (Section 4.3) has suggested an antagonistic interaction between powdery mildew and the yeast *Buckleyzyma aurantiaca* (Pauvert et al., 2020), which remains to be experimentally validated with culture-dependant approaches (Section 4.4). Moreover, as for *Plasmopara*

*viticola*, the virome of *E. necator* has been studied in search of hypovirulence-inducing mycoviruses (Pandey et al., 2018). Mycoviruses have been detected but an effect on fungal virulence and disease development has yet to be demonstrated.

Future studies should reinforce the searching for members of the resident microbiome and virome that naturally regulate powdery and downy mildews, and identify the biodiversity management practices that favour these natural antagonists. Such results are urgently needed to move towards pesticide-free viticulture (Jacquet et al., 2022), as mildews are currently controlled by chemical pesticides whose use should be drastically reduced to preserve environmental and human health. Based on the co-evolutionary history of *Vitis* spp. and mildew agents (Section 2.2), we recommended exploring the microbiome and the virome of mildew-resistant *Vitis* species in their native range (Fig. 2) to identify biocontrol candidates. In addition, we would recommend comparing the leaf microbiome of European cultivated grapevines that are rarely *vs.* frequently infected with mildews, according to epidemiological records (Section 4.1). The upper soil and bark microbiomes could also be analysed as they constitute the overwintering sites of the pathogens. Identifying the microbial taxa that are enriched in mildew-resistant *Vitis* species and in rarely-infected plots, and deciphering the microbial interaction networks (Section 4.3), could provide interesting insights into the natural regulation of mildews.

### 3.4 Dynamics of the berry microbiome to ripening and winemaking

Like the leaf microbiome (Section 3.3), the berry microbiome varies significantly in abundance and composition throughout the season. Population sizes of both fungal and bacterial communities on the berry surface increase with ripening and reach a maximum when the berries become overripe. The community composition varies significantly during the season, with a decrease in *Basidiomycota* (*Aureobasidium*, *Cryptococcus*, and *Rhodotorula* spp.) and an increase in *Ascomycota* (*Hanseniaspora*, *Metschnikowia*, *Pichia* spp.) (Barata et al., 2012). As the season progresses, Gram-negative bacteria (mostly *Pseudomonas* spp.) decrease in abundance whereas Gram-positive communities (mostly *Micrococcus* spp.) increase (Martins et al., 2012). The morphology and chemical composition of berries are major drivers of these seasonal variations in microbiome composition (Liu and Howell, 2021). Indeed, the water activity on grapevine berry skin and the composition of exudates change along the grapevine berry ripening process, with an

increase in sugar content, a decrease in pH and acidity (Griggs et al., 2021; Martins et al., 2014, 2020). Other drivers include agricultural practices during the fructification period (phytosanitary treatments, pruning), as well as the climatic conditions from fruit set to harvest (Belessi et al., 2022; Bokulich et al., 2014; Cordero-Bueso et al., 2011a,b; Ding et al., 2021; Martins et al., 2014, 2020; Pinto et al., 2014). In addition to these seasonal variations, the berry microbiome varies markedly in space. Over the last decade, many studies conducted at different spatial scales (world, countries, regions and vineyards) have provided evidence for a biogeographic differentiation of the berry microbiome (Bokulich et al., 2014; Gobbi et al., 2022; Jara et al., 2016; Li et al., 2022; Miura et al., 2017; Vitulo et al., 2019). Based on these findings, these authors have developed the concept of “microbial terroir” that proposes a causal relationship between grapevine microbial biogeography and regional wine characteristics.

At the ripening stage, the grapevine berries harbour fermentative microorganisms that will subsequently contribute to the winemaking process and to wine quality (Fig. 3). These microorganisms include *Saccharomyces cerevisiae* (Sections 2.1 and 2.3), non-*Saccharomyces* yeasts and Lactic Acid Bacteria (LAB). *S. cerevisiae* is the species most responsible for alcoholic fermentation. Although its populations reach their highest abundance at berry maturity ( $10^4$ – $10^6$  cfu/g), the yeast remains rare at the grapevine berry surface, thus raising questions about its origin (Börilin et al., 2020; Mortimer and Polsinelli, 1999). The abundance and diversity of fermentative yeasts on the grapevine berry surface is impacted by fungicide treatments (Cordero-Bueso et al., 2014) and infection by *Botrytis cinerea*, which modifies sugar availability (Fleet, 2003; Nisiotou and Nychas, 2007). Low numbers ( $<10^3$  cfu/g) of cultivable LAB are also present on sound fruit and end up in must during the early stages of wine processing. Within this group of bacteria, *Oenococcus oeni* (Fig. 3) is responsible for a crucial process called malo-lactic fermentation (MLF), where L-malic acid in wine is converted to softer L-lactic acid. The reduction of acidity, which spontaneously occurs after alcoholic fermentation, is beneficial to the quality of wines made in cool winegrowing regions. Yet, MLF only starts when the *O. oeni* population reaches  $10^6$  cfu/mL. Understanding the factors reducing the bacterial biomass is crucial, in order to avoid slow or blocked MLF. Phages (Fig. 3) are one of these factors, as they may modulate *O. oeni* population size, and a number of temperate and virulent phages infecting *O. oeni* have been recently isolated from crushed grapes and later winemaking stages (Claisse et al., 2021).

The berry microbiome may also contain microorganisms that alter wine quality, such as Acetic Acid Bacteria (AAB) or fungi that produce off-flavours or mycotoxins. AAB (*Acetobacter* spp., *Gluconobacter* spp.) (Fig. 3) are wine spoilage bacteria that are naturally present in the microbiome of grapevine berries (Fleet, 1993). They cause pre- and post-harvest yield losses and render berries unsuitable for wine production. A relevant example is sour rot of grape (Leixa et al., 2018), which is a polymicrobial disease requiring the presence of *Drosophila* flies (disease vector), wounded fruits, various yeasts that convert grape sugars to ethanol, and AAB that subsequently oxidise the ethanol to acetic acid. AAB co-exist with phages on ripe berries and a member of the *Tectiviridae* family has been recently characterised (Philippe et al., 2018). The control of AAB is challenging as these bacteria show great capacities to persist during wine making. Phages and their derived enzymes called endolysins may offer new solutions to their control.

Fungi naturally present on ripened grapevine berries can also produce off-flavours, in particular earthy or mouldy odours in grape juice and wines. This is the case of *Botrytis cinerea* (Fig. 3), the necrotrophic fungal pathogen responsible for grey mould (Armijo et al., 2016). *B. cinerea* produces cellulases, pectinases, proteases and a *p*-phenol oxygen oxidoreductase able to oxidise phenolics compounds, which is at the origin of severe alteration of wine colour. *B. cinerea* also causes an enhancement of phenylpropanoid metabolism, terpenes and fatty acid aroma precursors at the grapevine berry level. It can be associated with *Penicillium expansum*, then leading to the production of geosmin, an earthy off-aroma (La Guerche et al., 2005, 2007). However, *B. cinerea* can also colonise berries under its “noble rot” form which positively develops the berry aroma potential (Thibon et al., 2010, 2011). Natural antagonists of *B. cinerea* have been searched for within the leaf and berry microbiome to develop microbial biocontrol. They include *Aureobasidium pullulans*, which acts against *B. cinerea* through nutritive and spatial competition and antibiosis (Carmichael et al., 2019; Martini et al., 2009, p. 200; Rathnayake et al., 2018). They also include some bacteria (*Bacillus* spp., Actinomycetes) and yeasts (*Metchnikovia* spp., *Pichia* spp.) (Loqman et al., 2009; Raspor et al., 2010; Santos and Marquina, 2004). Mycoviruses infecting *B. cinerea* (Pearson and Bailey, 2013; Ruiz-Padilla et al., 2021) have been searched for to foster biocontrol. Although a rich virome was discovered, only a few mycoviruses were associated with a reduced virulence of the fungal pathogen (i.e., an hypovirulence phenotype) (Khalifa and MacDiarmid, 2021; Wu et al., 2010; Yu et al., 2015). These mycoviruses belonged to several families, including *Namnaviridae*

(single-stranded RNA virus) and *Genomoviridae* (single-stranded DNA virus). An unclassified double-stranded RNA virus was also found. Finally, fungal pathogens producing mycotoxins (*Aspergillus carbonarius*, *Aspergillus niger*) can also develop in mature berries. Mycotoxin contamination can be limited by specific practices (Hocking et al., 2007; Perera et al., 2021), including biocontrol by *A. pullulans* (Bozoudi and Tsaltas, 2018).

Grapevine berries constitute a pivotal microbial compartment because the dynamics of their microbiome depends on the agricultural practices and the climatic conditions in the vineyard, and determines the quality of the end product. Future research should therefore continue investigating the seasonal dynamics of microbial interactions and functions on the grapevine berry surface (Section 4.3), to elucidate their impact on berry quality and wine properties. Significant advances using metabolic networks (Section 4.3) were made for *Oenococcus oeni* and should be extended to the microbial community level, to understand the interactions between *O. oeni* and other microorganisms (like that *Saccharomyces cerevisiae*, LAB and AAB) that belong to the berry microbiome and are involved in winemaking (Mendoza et al., 2017). Conversely, the impact of the winemaking process on the berry microbiome dynamics could be assessed. Indeed, during winemaking, fermentative microorganisms reach very high population levels in cellars and can be disseminated in the vineyard through CO<sub>2</sub>-extraction systems and insect vectors.

### 3.5 Wood microbiome dysbiosis during grapevine aging and decline

Cultivated grapevines are perennial plants that age and are eventually removed when they are no longer productive. Nowadays, a major cause of grapevine decline (Fig. 3) are Grapevine Trunk Diseases (GTDs), a group of diseases which re-emerged in the late 1990s and cause wood degradation and necrosis within the inner structures of mature grapevines (Bertsch et al., 2013; Bruez et al., 2013; Mugnai et al., 1999). GTDs include *Botryosphaeria* dieback, *Diaporthe* dieback, *Eutypa* dieback, *Phomopsis* dieback, Petri disease, black-foot disease and Esca complex diseases. Pathogenic fungi associated with GTDs belong to several genera, including *Phaeoconiella*, *Phaeoacremonium*, *Fomitiporia*, *Eutypa*, and to the *Botryosphaeriaceae* family (Cobos et al., 2022; Fig. 3). However, these fungi have been isolated from both healthy and necrotic wood tissues of both GTD-symptomatic and asymptomatic grapevines. Their role in GTDs has therefore been questioned

(Hofstetter et al., 2012) and other microorganisms potentially involved in GTDs have been isolated using culture-dependent methods (Elena et al., 2018; Kraus et al., 2019). Cobos et al. (2022) reported that in 2018, 133 fungal species belonging to 34 genera (mainly ascomycetes and basidiomycetes) had been associated with GTDs worldwide, this number increasing yearly. Bacteria inhabiting grapevine wood have also been isolated, identified and their functions studied. For instance, Bruez et al. (2015) identified 26 bacterial genera inhabiting the various wood tissues, the most abundant being *Bacillus* (34% of the bacterial strains), then *Pantoea* (12%), *Paenibacillus* (9%), and *Enterobacter* (6%). Bacterial communities differed between anatomical parts (i.e., trunk or cordon) and tissue types (i.e., necrotic or non-necrotic). Functions of bacteria colonising necrotic and non-necrotic tissues differed, as they metabolised differently the carbon substrates. Over the last decade, next-generation sequencing (Del Frari et al., 2019; Eichmeier et al., 2018; Gramaje et al., 2022; Lade et al., 2022) and metatranscriptomic analyses (Nerva et al., 2022; Paolinelli et al., 2022) have provided a deeper resolution in the grapevine wood fungal and bacterial microbiomes. Depending on the grapevine organs, years, seasons and wood health status (i.e., healthy or necrotic), variations in the wood microbiome composition were observed, for both fungi and bacteria (Bruez et al., 2014, 2020).

Wood necrosis is often observed in mature grapevines (15–25-year-old), the difference between healthy and GTDs-plants being the quantity of wood necrosis. White-rot necrotic wood, in particular, is more abundant in diseased plants (Maher et al., 2012; Ouadi et al., 2019, 2021). This observation raised a key question: why does a substantial percentage of wood become necrotic in mature plants? This question has recently been investigated in the case of Esca, the most frequent GTD. Microbiome changes occurring at the onset of Esca were investigated using metabarcoding approaches (Bruez et al., 2020). The results showed a decrease in the diversity of fungal communities in the white-rot of diseased trunks, linked with the tissue colonisation by the fungal pathogen *Fomitiporia mediterranea*. Because fungi and bacteria coexist in the wood, the authors hypothesised that fungal-bacteria associations were involved in the wood degradation. In agreement with this hypothesis, associations between GTD-fungal pathogens (*F. mediterranea* and *Phaeoconiella chlamydospora*) and bacterial species (*Sphingomonas* spp. and *Mycobacterium* spp.) were detected in the cordons of young grapevines expressing Esca-foliar symptoms. More recently, Haidar et al. (2021) showed that *F. mediterranea* and a novel species of bacteria isolated from



grapevine wood (Haidar et al., submitted) acted synergistically to enhance wood structure degradations associated with Esca.

Plant-microbe and microbe-microbe interactions responsible for other GTDs (*Botryosphaeria dieback*, *Diaporthe dieback*) have also been studied. Several fungal pathogens of the genera *Botryosphaeria*, *Diplodia*, *Lasiodiplodia* and *Neofusicoccum* have been identified as causal agents of *Botryosphaeria dieback*. This complex of xylem-inhabiting fungi can remain latent in the roots before producing phytotoxic compounds and degrading enzymes (xylanases, cellulases,  $\beta$ -glucanases) in stems and symptomatic leaves (del Martínez-Diz et al., 2019; Gramaje et al., 2018; Marais et al., 2021). Fungi of the genus *Diaporthe* are responsible for *Diaporthe dieback*, a disease that causes, among others, cane bleaching, swelling arms, trunk cankers, shoots breaking off at the base, loss of vigour, and reduced bunch size (del Martínez-Diz et al., 2019; Guarnaccia et al., 2018). These symptoms can allow the implantation of opportunistic microorganisms such as *Didymella negriana*, which is capable of producing mycotoxins (Liu and Howell, 2021; Stranska et al., 2022).

Wood bacteria, fungi and mycoviruses with potential biocontrol activities against fungal pathogens associated GTDs have been investigated (Haidar et al., 2021; Marais et al., 2021; Mondello et al., 2018; Rezgui et al., 2016). Bekris et al. (2021) analysed the wood microbiome of asymptomatic and symptomatic plants in three Greek grapevine varieties, using differential abundance analysis (Section 4.1) and network inference methods (Section 4.3). Their results showed that *Bacillus* and *Streptomyces* genera dominated the bacterial microbiome of asymptomatic plants and had a negative co-occurrence pattern with GTD-pathogens belonging to the *Phaeoconiella*, *Phaeoacremonium* and *Seimatosporium* genera. The authors concluded that these bacterial taxa may play a role in the suppression of GTDs. The suppressive role of the wood microbiome was also suggested by Bruetz et al. (2016), as they noticed that pathogenic and non-pathogenic fungi reach an equilibrium in the functional tissues of old grapevines. Mycoviruses may also play a role in the suppression of GTDs. For instance, mycoviruses were detected in *Neofusicoccum parvum* isolates from grapevines without symptoms of Esca (Nerva et al., 2019) and in hypovirulent isolates (Marais et al., 2021), suggesting that mycoviruses may reduce the aggressiveness of this *Botryosphaeriaceae* species. It is noteworthy that viruses could also be helpful to fight pathogenic bacteria that colonise xylem vessels, like *Xylella fastidiosa* (Fig. 3), the causal agent of Pierce's Disease, and

*Xylophilus ampelinus*, the causal agent of bacterial blight. Biocontrol protocols using virulent phages have been successfully implemented *in planta* in Texas, USA, to control *Xylella fastidiosa* (Das et al., 2015). This approach is promising, especially since the pathogen is vectored by an insect (the glassy-winged sharpshooter, *Homalodisca vitripennis*) (Bhowmick et al., 2016) which has the capacity to uptake *X. fastidiosa* phages (Clavijo-Coppens et al., 2021).

Deciphering plant-microbe-virus interactions occurring in the grapevine wood is crucial for future monitoring and management of GTDs. Specifically, tipping points in the wood microbiome dynamics related with the onset of disease symptoms could inform disease-monitoring programmes (Section 4.3). For future research, we would recommend analysing the taxonomy and functions of the wood microbiome, as well as the changes in microbial interactions networks (Section 4.3), just before the appearance of the first Esca-foliar symptoms to answer the following questions: (i) is the onset of the disease marked by an increase in the heterogeneity of the microbiome between grapevines, as stated in the “Anna Karenina principle” (Zaneveld et al., 2017)? (ii) is the onset of the disease marked by a collapse in microbial interaction networks? (iii) is there a change in the functional composition of the microbiome, marked by an increase in the abundance of pathogenic species?

Answering these questions about the wood microbiome and those raised earlier for the other microbial compartments (roots, leaves, and berries) will require the study of the complex microbial communities associated with cultivated grapevines, to identify the environmental drivers of plant-microbe-virus interactions. It will also require the analysis of simpler, synthetic microbial consortia in controlled conditions, to get mechanistic insights into microbial interactions. Both approaches will be linked. Studying natural communities will require dedicated field sampling designs to identify key taxons and functions associated with grapevine performance. These key taxa and functions will then help design culture-based experiments coupled with modelling approaches to leverage omics data and decipher microbial interactions at the genome scale. These models will be used to optimise the microbial consortia (Busby et al., 2017; Toju et al., 2018) that could ultimately be inoculated to foster grapevine performance under conditions of climate change and reduced pesticide use. Some of the tools to address these challenges and make the grapevine microbiome more functional are described in the next section (Section 4).



## 4. Let's make the grapevine microbiome more functional

This section proposes experimental and computational approaches that could help uncover the functioning of the grapevine microbiome, with the goal of integrating it into a more sustainable viticulture. We propose field sampling designs and statistical approaches that exploit what we know of the evolution and dynamics of the grapevine microbiome (Sections 2 and 3) and aim to identify microbial taxa or consortia that may play a role in vine performance and vineyard ecosystem health. We then review the molecular tools and computational approaches that work together to analyse the grapevine holobiont as a dynamic, interactive, and functional system. We emphasise throughout the critical need for culture-dependent approaches to validate the effects of specific microbial taxa and functions on vine performance and vineyard ecosystem functioning. Each subsection sets out research hypotheses or questions about microbiome functioning, describes methodological frameworks that exist or could be developed to address these questions, and finally suggests how to apply them to the grapevine system.


### 4.1 Field sampling designs and statistical approaches to identify beneficial microbial taxa

Grapevine hosts many hundreds of microbial taxa (Knapp et al., 2021; Morgan et al., 2017; Zarraonaindia et al., 2015 and Section 3) of which only a few dozen have known roles and modes of action (Fig. 3). Here we propose field sampling designs and statistical approaches that could be applied to metabarcoding datasets to elucidate the links between the presence or abundance of particular microbial taxa, and functional traits measured at the plant- or ecosystem-scale. In the current context of climate change and need to limit pesticide use (Jacquet et al., 2022), we ask which field sampling designs and statistical approaches are suitable for identifying microorganisms enhancing plant tolerance to drought or providing a barrier effect against microbial pathogens?

#### ***Question #1: How to identify microbial taxa enhancing plant tolerance to drought or providing a barrier effect against microbial pathogens from metabarcoding data?***

To address this question, the composition of grapevine microbiome samples can be analysed using metabarcoding approaches and then statistically related

to measurements of grapevine physiology or vineyard ecosystem functioning. For example, the microbiome can be related to ecophysiological traits that indicate grapevine tolerance to drought (Gambetta et al., 2020; Vacher et al., 2022), or to the frequency and intensity of diseases at the plot level in epidemiological surveys (Chen et al., 2019), or to more integrative agricultural variables such as yield potential. The grapevine microbiome can then be considered as an explanatory factor of the variation in the functional trait of interest (e.g., drought tolerance, disease resistance, yield). This variation can be analysed at different temporal and spatial scales (Fig. 4): within



**Fig. 4** Workflow, from field experiments to mathematical models of omics data, to decipher the complex plant-microbe-microbe interactions forming the grapevine holobiont and to identify beneficial microbial taxa. The experimental design sets up the sampling plots, hosts, compartments and time-points, according to spatial and temporal variations in a functional trait of interest (e.g., grapevine drought tolerance, disease resistance, yield). Grapevine samples (roots, trunk, leaves or berries) are then collected, cultivated, amplified and sequenced, and omics data are processed. Omics approaches can be culture-based (culturomics), taxon-based (metabarcoding), gene-based (microfluidic qPCR chips, for instance) or function-based (shotgun metagenomics). Bioinformatic analysis and data integration pipelines result in multi-omics tables, time series, SynComs or GSMN construction. This data can be used as input data for interaction network inference, species or community-level models and dynamical systems representing microbial populations' dynamics under various environmental conditions. ASV, Amplicon Sequence Variant; GSMN, Genome-Scale Metabolic Network; MAG, Metagenome-Assembled Genomes; OTU, Operational Taxonomic Unit; SynCom, Synthetic Community.

the lifetime of a cultivated vine individual (Fig. 3), between vines planted in the same plot (e.g., Darriaut et al., 2021), between vineyard plots, between cultivated vines and their wild ancestors (Section 2.1), or between cultivated vines and disease-resistant *Vitis* species (Section 2.2).

The field sampling design should determine the statistical analyses to be used. One sampling approach is to examine grapevine individuals or vineyard plots along a gradient of a previously measured functional trait of interest. Another would sample pairs of individuals or pairs of plots, of contrasted functional traits (e.g., drought-tolerant vs. drought-sensitive, disease-resistant vs. disease-susceptible, high- vs. low-yielding), other effects being otherwise controlled (same rootstock and scion cultivars, similar plant age, soil type and farming system). These two sampling strategies (Fig. 4) rely on spatial variation in the functional trait of interest. The first maximises the variation in the trait along the gradient, while the second reduces the environmental heterogeneity within a pair. Longitudinal, time-series surveys (Fig. 4) would complete this picture, and highlight joint temporal variations between the microbiome and the functional trait of interest. This combination is highly relevant for grapevine given the strong dynamics in the microbiome at multiple timescales (Sections 2 and 3).

Several statistical tools can then be applied to metabarcoding data to assess the contribution of microbiome composition to variation in a functional trait, and identify the properties of the microbiome that have explanatory power. These explanatory properties can be the occurrence or abundance of particular microbial taxa or microbial functional groups (Section 4.2), or aggregated properties at the community or network levels such as community alpha-diversity or microbial network connectance (Barroso-Bergadà et al., 2021). The compositional dissimilarity matrix between samples (community beta-diversity) can be converted into explanatory vectors using the method of PCNM eigenvectors, developed for spatial distance matrices (Borcard and Legendre, 2002). The statistical model will have the functional trait of interest as the dependent variable and the microbiome properties and environmental factors as explanatory variables (e.g., Asad et al., 2022 for yield; Cambon et al., 2022 for drought tolerance; Pérez-Valera et al., 2020 for an example of paired samples). When the microbiome contributes significantly to the functional trait of interest, identifying which microbial taxa are the main players can be done using the TITAN2 method (Baker et al., 2015) or differential abundance analysis methods (Nearing et al., 2022; Weiss et al., 2017). Machine learning or deep learning methods can also be applied (Xu et al., 2022); for example, a Random Forest

algorithm has been used to identify combinations of microbial taxa indicative of yield (Yergeau et al., 2020). For grapevine, statistical identification of subsets of microbial taxa potentially involved in disease resistance or drought tolerance would be relevant to set up experimental, culture-dependant approaches (Sections 4.4). Finally, to test specific hypotheses on the direct and indirect effects of the explanatory factors (including microbiome) on the functional trait of interest, structural equation models (SEM) have been developed (Jassey et al., 2018). For grapevine, a descriptive SEM would be highly relevant for assessing the direct and indirect effects (through the microbiome) of variety and environmental factors (drought, agricultural practices) on disease severity and frequency.

## 4.2 Molecular tools to uncover the functional potential of the microbiome

Microbiome samples collected from contrasted environments following the field sampling designs described above (Section 4.1) can be further analysed using culture-independent molecular tools to uncover the microbial genomes present in the sample, analyse the microbial functions they carry, estimate precisely variation in their abundance and link this variation with plant gene expression patterns. Here, we review the molecular tools that could be used to address the following questions: How to characterise the functional potential of the grapevine microbiome with shotgun metagenomics; and, how to decipher the functional coupling between grapevine and its microbiome with quantitative approaches?

### ***Question #2: How to characterise the functional potential of the grapevine microbiome with shotgun metagenomics?***

Metataxonomics using metabarcoding of amplified marker DNA sequences has been widely used to inventory the microbial taxa present in a sample (Marchesi and Ravel, 2015), including in grapevine (Sections 2 and 3). Bioinformatic tools exploit this data to generate abundance tables of microbial Operational Taxonomic Units (OTUs) or Amplicon Sequence Variants (ASVs) in samples (Callahan et al., 2016, 2017; Estaki et al., 2020; Hildebrand et al., 2014; Pauvert et al., 2019; Schloss et al., 2009). Metabarcoding has strong limitations, however, including primer amplification bias, copy number variation of the target sequence across taxa, and limits to taxonomic resolution, which may prevent the accurate characterisation of taxa composition and abundance. Although primarily dedicated to taxonomic characterisation, metabarcoding data can nevertheless be used to

get insight into the functional potential of a microbiome with taxonomy-based function prediction methods (Belcour et al., 2022; Douglas et al., 2020; Louca et al., 2016; Nguyen et al., 2016; Wemheuer et al., 2020; Zanne et al., 2020). These methods build upon the association of taxonomic clades to known functional guilds, or to metabolic functions or pathways obtained from public databases of annotated genomes (Section 4.3).

In contrast to metabarcoding approaches, shotgun metagenomics (Fig. 4) does not sequence DNA markers but the total DNA material of a sample. Significant data processing is required (Bickhart et al., 2022). Reference-based profiling (Section 4.3) compares these reads to reference genomes. In genome-resolved metagenomics, quality-filtered reads are assembled into contigs (Li et al., 2015; Nurk et al., 2017), which can be parsed into genes that are binned into metagenome-assembled genomes (MAGs) (Borderes et al., 2021; Kang et al., 2019; Wu et al., 2016; Yue et al., 2020). Genomes can be resolved for uncultured bacterial and eukaryotic members of the microbiome, and the precise function carried by a species in the sample can be investigated. The limitations of MAG reconstruction come about as a consequence of the diversity of species found in the same sample (Lemos et al., 2021; Nayfach et al., 2020). Improvements to long-read sequencing such as PacBio or Nanopore technologies lead to better metagenome assembly. They can be combined synergistically with low error rate short reads to accurately retrieve repetitive genome regions (Chen et al., 2021; Damme et al., 2021). Long reads using the most recent technologies can however suffice alone for reconstructing high-quality MAGs (Sereika et al., 2022). The principal bias in shotgun metagenomics is the taxonomic and functional incompleteness in reference databases that can add noise to the taxonomic assignment and the functional annotation for underrepresented species and functions. In addition to shotgun metagenomics, metabolomics and metatranscriptomics (Paolinelli et al., 2022) approaches identify metabolites or expressed genes, respectively, in the host and the microbiome.

While genome-resolved metagenomics applications to soil and plants are becoming more frequent (Wu et al., 2022a,b), few studies have yet applied it to grapevine. Metagenomics has been used to analyse the microbial composition of grapevine berries and to reconstruct 13 near-complete draft genomes (Salveti et al., 2016). Azevedo-Silva et al. (2021) discussed the feasibility of applying shotgun metagenomics to grapevine leaf, rhizosphere and soil characterisation and proposed a protocol. For grapevine, we recommend using metataxonomics before shotgun metagenomics to first target



the environmental samples worthy of deeper sequencing (Fig. 4). We argue that, while metagenomics has some biases, it has fewer than amplicon sequencing and the substantial amount of information it brings makes it worthy of consideration for deeper analyses of the grapevine microbiome.

***Question #3: How to decipher the functional coupling between grapevine and its microbiome with quantitative approaches?***

Metabarcoding and metagenomics data are by nature compositional, meaning that we measure the relative abundance of microbial taxa or functions within a sample rather than their absolute abundance. To produce absolute counts requires complementary, quantitative methodologies (Props et al., 2017). Real-time quantitative PCR (RT-qPCR) has developed into a powerful and accurate technique for measuring absolute abundance by assessing quantitative changes in taxonomic or phylogenetic marker genes across spatial and temporal scales (Heid et al., 1996), supplanting the classical colony counting method, which is hindered by the low proportion of cultivable microorganisms (Lou et al., 2018). Development of automated, high-throughput qPCR platforms (Fig. 4) have increased, 10-fold, the number of qPCR reactions that can be simultaneously performed on a single instrument. Microfluidic qPCR chips with the Biomark<sup>®</sup> HD system, for example, can perform 9216 qPCR reactions (96 samples amplified with 96 primer sets) in only 4h.

Different microfluidic qPCR chips have been used in pioneer studies of gene expression in grapevines. The NeoVigen chip (Dufour et al., 2016), developed to quantify defence gene expressions in grapevine, quantifies plant genes involved in pathogenesis-related protein synthesis, secondary metabolites synthesis, the indole pathway, the oxidoreduction system, the salicylic acid, ethylene, and jasmonic acid pathways, and cell wall reinforcement. The BioStim chip (Bodin et al., 2020) has extended the gene set by including, among others, genes involved in primary pathways and phytohormones pathways (auxin, cytokinin, gibberellin, abscisic acid). These chips have been used to decipher the mechanisms underlying the effects of biostimulants on grapevine resistance to multiple stresses. The workflow can be adapted to different types of microbial marker genes, whether taxonomic (Dreier et al., 2021; Kleyer et al., 2017) or functional (Crane et al., 2018). The technology could therefore provide simultaneous quantitative snapshots of the microbiome taxonomic and functional composition and of the host response (Noman et al., 2021; Zolti et al., 2020).

To decipher the functional coupling between the grapevine host and its microbiome, we recommend parallel development of microfluidic chips targeting plant genes involved in plant growth or stress resistance, and chips targeting specific microbial taxa and functions. Considerable progress has been made for the plants (Bodin et al., 2020; Dufour et al., 2016), but a chip has yet to be created for microbiomes. Several challenges must be overcome to choose the most relevant microbial targets, design primer sets, and validate the chip. Target selection must deal with the multitude of microbial taxa colonising the plant and the multitude of functions they carry (Section 3). Chip validation must use reference species and synthetic microbial communities (Section 4.4) before applying the chip to environmental samples. Once developed, this tool could link microorganisms to plant health biomarkers, for instance to identify beneficial (phyto-probiotic) microbial agents that foster plant health or stimulate plant immunity, or potential antagonists of vine pathogens that are candidates for next-generation biocontrol products.

### **4.3 Computational approaches to understand the grapevine holobiont as a functional and dynamic network**

Computational methods exploit the wealth of -omics data produced by high-throughput methods (Sections 4.2) to infer networks of ecological interactions between microbial taxa, unravel the trophic and metabolic relationships that underlie them, and analyse how they drive microbiome dynamics at different time scales (Sections 2 and 3). The various types of -omics data (metataxonomics, shotgun metagenomics, metranscriptomics, metaproteomics, metabolomics) are complementary and can be used to reconstruct ecological networks, metabolic networks, protein-protein interaction networks and gene regulatory networks (Liu et al., 2021). Here we describe the tools to specifically answer the following questions: How to characterise ecological interactions between grapevine-associated microorganisms using metabarcoding data? How to decipher metabolic interactions within the grapevine microbiome using shotgun metagenomics? How to infer the microbial interactions that shape microbiome dynamics from time-series data?

#### ***Question #4: How to identify and characterise ecological interactions between grapevine-associated microorganisms using metabarcoding data?***

Microbial ecologists have long used metabarcoding to learn ecological interactions between microbial taxa. They assume that since past or ongoing

interactions between microbial taxa may impact their frequencies, patterns in relative shifts in frequency may provide information about the structure of the ecological interaction network (Cobo-Díaz et al., 2022; Jakuschkin et al., 2016; Kerdraon et al., 2019; Poudel et al., 2016; Vacher et al., 2016b). However, this process is not straightforward: metabarcoding data can be biased by amplification and sequencing artefacts, and are always compositional (Gloor et al., 2017; Section 4.2). One of the first tools for inferring interaction networks from relative sequence counts was SparCC (Friedman and Alm, 2012), which avoided the compositional effect by using log-transformation components to obtain correlation. Other tools like SPIEC-EASI (Kurtz et al., 2015) or CCLasso (Fang et al., 2015) refined correlation-based measures. PLN (Chiquet et al., 2019) enhances inference accuracy by exploiting sample covariance, while MPLasso (Lo and Marculescu, 2017) introduces the use of prior microbiological knowledge through data-mining of external sources of information. HMSC (Ovaskainen et al., 2017) also integrates covariates on samples and taxa and was recently used to infer microbial networks (Abrego et al., 2020; Fort et al., 2021). These tools are extremely robust to noise in experimental data and can be run rapidly on even quite large datasets. Most of them are available in R, which facilitates their use (Dohlgan and Shen, 2019; Tikhonov et al., 2020). However, these tools define no clear link between the positive and negative interactions they find, and the ecological mechanisms that may generate them (Röttgers and Faust, 2018).

Explainable machine learning can relate ecological mechanisms to inferred interaction networks (Tamaddoni-Nezhad et al., 2021). Different interaction types, such as competition and mutualism, lead to different changes in species frequencies and patterns (Fig. 4), which can be learned (Derocles et al., 2018; Tshikantwa et al., 2018). Explainable machine learning transcribes these criteria into logical rules that directly infer ecological interactions, with little to no human intervention in the interpretation step (Barroso-Bergada et al., 2022). Hypotheses of ecological interactions between taxa pairs are generated through a process that simultaneously detects interactions and classifies them as competition, mutualism, amensalism or commensalism. This direct approach is particularly valuable for reconstructing microbial networks in previously unstudied ecosystems, where human knowledge for interpretation may effectively be lacking.

For grapevine, this approach was recently applied to microbial communities sampled from symptomatic and asymptomatic leaves of European cultivated grapevine during downy mildew epidemics. After a thorough assessment of the method using simulated datasets (Weiss et al., 2016),

preliminary interaction networks were computed and showed potential antagonists of the causal agent *Plasmopara viticola*—some known, some new—constituting options to control mildew development. We expect that this approach will identify sub-networks of interest, centred on pathogen species and their direct antagonists or facilitators, whose subsequent modelling could lead to better understanding of pathogen-microbiome interaction dynamics.

### **Question #5: How to decipher metabolic interactions within the grapevine microbiome using shotgun metagenomics?**

The microbial functions present in a given sample can be assessed from shotgun metagenomic surveys or from metataxonomic data after taxonomy-based function prediction (Section 4.2). An advantage of the former is functional profiling of communities, using assembly-independent reference-based approaches, or de-novo reference-free assembled sequences (Quince et al., 2017). Both approaches delineate microbial species and strains, needed for reconstructing microbial metabolic networks. These networks are then used in mathematical models (kinetic, constraint-based and discrete models) to predict metabolic reactions at the species- or community-level.

Functional profiling of microbial communities using assembly-independent reference-based approaches directly compares reads to reference databases (Beghini et al., 2021) to identify genes and thereby functions. Functions or pathways associated with samples are then compared or used in statistical learning in order to reveal higher level functional patterns between sample groups (Raguideau et al., 2016). Unfortunately, reads cannot be linked to their genome of origin, impeding the interpretation of these patterns in terms of metabolite-mediated trophic interactions between microbial taxa. Moreover, plant and grapevine-associated reference catalogues are less available than, for instance, the gut microbiome. Hence, genome-resolved metagenomics, below, seems to be a better option for grapevine.

Reference-free de novo assembly allows functional annotation of genes (Cantalapiedra et al., 2021; Seemann, 2014), functional characterisation of the corresponding organisms, and genome-scale metabolic network (GSMN) reconstructions (Fig. 4). GSMN consist of all biochemical and transport reactions predicted from an individual microbial genome (Gu et al., 2019) and can be reconstructed by different tools (Arkin et al., 2018; Karp et al., 2019; Seaver et al., 2021) or even directly connected to MAG reconstruction (Zorrilla et al., 2021). In most cases,

GSMN reconstruction involves manual curation to ensure the quality of the predictions they provide (Karp et al., 2018). GSMN can be associated with mathematical models to predict the metabolism of a species or a community of organisms in an environment. The main difficulty is to achieve high-quality GSMNs, since (i) MAGs are frequently incomplete, even worse so when under-studied microbiota are under-represented in the gene annotation databases; and (ii) GSMN reconstruction from the MAG can be imperfect. Manually curated GSMNs are best but the number of genomes in a microbiota and limited knowledge about the microorganisms argues in favour of automatisisation (Belcour et al., 2020). Fortunately, the multiplication of shotgun sequencing studies of the grapevine microbiome will increase the size and quality of grapevine-related annotated gene catalogues and favour better MAGs-based GSMN automatic reconstruction

Kinetic modelling uses differential equations models of the kinetics of biochemical reactions throughout the metabolic network of an organism (Ross and McMeekin, 1994), permitting microbial phenotype prediction that accounts for enzyme and metabolite concentration dynamics. While it has been applied to model organisms such as *Escherichia coli*, it is not widely applied due to the need to obtain biochemical measurements to parameterize the models (Khodayari et al., 2014). An alternative are stoichiometric models, also known as constraint-based models. They translate a GSMN into a linear system of constraints for an optimisation problem used to predict metabolic fluxes across the GSMN under simplifactory assumptions such as steady state in the system, i.e., the non-accumulation of internal metabolites and their constant concentration over time. Flux Balance Analysis (FBA) or Flux Variability Analysis (FVA) compute the activity rates of metabolic reactions while optimising an objective function, typically the maximisation of the microbial population growth (Orth et al., 2010). FBA models can be coupled with differential equations through dynamic FBA (dFBA) (Mahadevan et al., 2002). These methods have been widely used and extended to the study of microbial communities (Biggs et al., 2015; Frioux et al., 2020) and to study the metabolism of oenological yeasts (Eder et al., 2020; Nidelet et al., 2016).

Discrete models are an alternative to constraint-based models to capture the metabolic potential of a GSMN, in a nutritional environment ideally characterised by metabolomics. Network expansion (Ebenhöh et al., 2004) computes the set of reachable metabolites from a given set of metabolic seeds and has been used as a building block for screening the metabolism of microbial communities (Belcour et al., 2020). Complexity

is reduced by selecting minimal communities that satisfy a metabolic objective, which may take the host metabolism into account (Frioux et al., 2018). Hypotheses arising from these minimal communities can be tested using synthetic communities (SynComs; Marín et al., 2021) selected to provide a targeted function. These approaches can be coupled with dFBA: discrete models are used for systematic screening and selection of SynComs, then dFBA is used on the selected SynComs to decipher the dynamics of the community at a molecular level.

GSMN were used to study soil and rhizosphere microbiome (Jacoby et al., 2017), and to model winemaking process (Mendoza et al., 2017). However, to our knowledge, GSMNs of grapevine phyllosphere or berry microbiomes remain an open challenge, of great potential for improving functional characterisation of microbiomes. For grapevine, kinetic modelling would permit, for example, integration of multi-omics data to decipher and validate the dynamical interactions between a microbial pathogen and putative antagonists. Minimal communities identified by discrete modelling, for example, to compete against microbial grapevine pathogens, or to promote oenological yeast growth by maintaining a tailored nutritional niche, can be tested with SynComs (Section 4.4).

### ***Question #6: How to infer microbial interactions shaping microbiome dynamics from time-series?***

Four fundamental, intrinsically dynamical ecological processes drive microbial communities: evolutionary diversification, selection, dispersal, and ecological drift (Nemergut et al., 2013; Vellend, 2010). The contribution of these deterministic and stochastic processes, complementary to historical contingency and priority effects, can now be evaluated cost-effectively using -omics time-series data (Ning et al., 2019; Vass et al., 2020; Zhou and Ning, 2017). Longitudinal time-series data enable the study of the dynamical interactions that determine selection in the microbial ecosystem.

One increasingly common approach is the use of Generalised Lotka-Volterra models (GLV) to decipher the dynamic interactions shaping community dynamics (Faust et al., 2018). GLV models decompose microbial growth into an intrinsic growth rate that models population fitness and interspecies interactions modelling ecological processes such as competition or parasitism (Fig. 4). These interactions are fitted to reproduce temporal data, which permits an ecological interpretation of the community dynamics. Despite simplifying assumptions such as constant environmental constraints, GLV models are suitable for studying microbiome stability,

microbial population persistence and microbial community robustness (Gonze et al., 2018). They model dynamics that correlation-based methods cannot (Carr et al., 2019). They have been used to study the dynamics of the gut microbiome during primary succession (Marino et al., 2014) and pathogenic infection (Stein et al., 2013), and to identify the assembly rules that determine the long-term composition of synthetic communities (Friedman et al., 2017). Computational tools for inferring GLV interactions have been proposed by Bucci et al. (2016) and Kuntal et al. (2019).

GLV models need specific experimental design to guarantee accuracy of microbial network inference (Remien et al., 2021). GLV models describe the evolution of absolute population, so model parametrization requires time-series of absolute counts (e.g., qPCR or flux cytometry, Section 4.2). Abundance proxies can be used, such as compositional metabarcoding data renormalized in each sample by the dosage of total DNA (Stein et al., 2013). The need for absolute population counts has been relaxed with an alternative GLV formulation for compositional data (Joseph et al., 2020). GLV models are better suited to analyse marked changes of the microbial community, for example, after strong perturbations such as pesticide treatment or colonisation by exogenous microbes such as pathogens (Remien et al., 2021), although Xiao et al. (2017) does address the steady-state microbiome.

GLV models have intrinsic limitations. First, growth and interspecies interactions are treated phenomenologically without regard to their underlying mechanisms, contrary to other models that supplement pairwise ecological interactions with metabolite-mediated interactions (Momeni et al., 2017). Second, microbial populations are treated as well-mixed without regard to spatial structures caused by nutritional niches (Prosser et al., 2007), which can impair GLV estimates (Armitage and Jones, 2019). Other population dynamics modelling frameworks avoid these drawbacks: GSMN, above, are particularly suitable for describing the metabolism of species subject to trophic interactions such as competition for resources, or cross-feeding in small consortia (Widder et al., 2016); spatially explicit microbial population dynamics models can take into account spatial heterogeneities (Labarthe et al., 2019). These models require, for parameterization, experimental measurements of reduced synthetic communities in controlled environments.

For grapevine, GLV models have not yet been used, but have been to model microbial dynamics in other plant species (Becerra-Lucio et al., 2021). GLV models could be coupled with machine-learning approaches



to select pathogen-associated sub-networks before studying their dynamics. They could also be used to study the dynamics of minimal communities identified by discrete modelling.

#### 4.4 Culture-dependent approaches to validate microbial interactions and functions

All statistical and computational approaches described above generate hypotheses about the microbiome and its dynamics that must be experimentally validated. Therefore, we ask, how to isolate and culture microorganisms from a microbiome to study their functions and their interactions?

##### ***Question #7: How to isolate and culture microorganisms from the grapevine microbiome to study their functions and their interactions?***

Culturomics, a high-throughput combination of multiple culture conditions and rapid identification of microorganisms (Fig. 4), has revitalised culture-dependent methods. The landmark studies by Lagier et al. (2015, 2016, 2018) combined high-throughput cultivation with cost-effective rapid identification by Matrix-Assisted Laser Desorption/Ionisation Time Of Flight Mass Spectrometry (MALDI-TOF MS), and increased the number of microorganisms identified in the human gut microbiome by more than 50%. Large-scale isolation strategies have since been developed, including new culture media and advanced cultivation technologies. Specific protocols consisting of successive dilutions of a microbial population to the point of no growth, and technologies such as microdroplet and microfluidics, have been proposed to enhance isolation efficiency (Button et al., 1993; Kaminski et al., 2016; Zhang et al., 2021). For the plant microbiome, plant-based media that isolate microorganisms in environmental and nutritional conditions close to their natural habitats have been successfully developed to improve cultivability (Mourad et al., 2018; Sarhan et al., 2018, 2019).

The success of culturomics in improving the catalogue of isolated human bacteria and plant endophytic bacterial communities (Bilen et al., 2018; Riva et al., 2022) holds promise for the grapevine microbiome, but the task ahead is huge. Fewer than 3000 plant-associated bacterial genomes are available, out of 70,000 known bacterial genomes (Joint Genome Institute Integrated Microbial Genomes; <https://img.jgi.doe.gov/>; Trivedi et al., 2021). High-quality sequencing, assembly, and annotation of pure microbial strain genomes is needed to improve gene and protein databases and

knowledge of the genome-based mechanisms underlying plant-microbe interactions (Levy et al., 2018; Sarhan et al., 2019; Section 4.3). Genome-resolved metagenomics produces uncurated draft genomes whose gaps, assembly errors and contamination from other genomes limit their usefulness (Chen et al., 2020); obtaining reference bacterial genome sequences (Trivedi et al., 2021; Section 4.3) will improve quality. Microbial culture collections make it possible to screen for microbial phenotypic traits that foster plant growth (growth promoting bacteria) and plant health (biocontrol agents) and they are imperative for validation of candidate genes. Cultivation and co-cultivation strategies are needed for experimental validation of predictive models of plant-microbiome and pathogen-microbiome interactions, and to provide deeper insight into the functional potential of beneficial plant-associated microbes (Nai and Meyer, 2018; Trivedi et al., 2020).



## 5. Conclusion and perspectives

The grapevine microbiome has developed over thousands of years, integrating taxa from Asia, Europe and America, under the pressures of human selection and pathogen invasions. There are many gaps in this history, but the numerous studies of the grapevine microbiome show that it contains beneficial microorganisms (plant growth-promoting bacteria, mycorrhizae, biocontrol agents, microorganisms of oenological interest) whose mode of action is sometimes well documented. These beneficial microorganisms belong to a dynamic microbiome that significantly varies throughout the plant lifetime and throughout the year. The initial microbiome of grafted plants is determined by the microbiomes of mother plants that produce rootstocks and scions. Those initial microbiomes may contain phytoviruses and fungal pathogens, but also beneficial microorganisms increasing tolerance to biotic and abiotic stressors. Some of the beneficial microorganisms (for instance, the arbuscular mycorrhizal fungi *Rhizophagus irregularis*) can be inoculated at the nursery stage to strengthen the microbiome. Once planted, roots actively release exudates that attract the rhizosphere microbiome from the soil reservoir. Beneficial microorganisms recruited by grapevine roots include plant-growth promoting bacteria (e.g., *Pseudomonas* spp., *Rhizobia* spp.) that make mineral nutrients (phosphorus, nitrogen and phosphate) available to the plant, arbuscular mycorrhizal fungi (e.g., *Glomus* spp.) that help the plant to cope with water

stress, and microorganisms whose biocontrol activity (e.g., *Bacillus* spp., *Pythium oligandrum*) relies on various types of microbial interactions including antibiosis, hyperparasitism, spatial and nutritional competition and interference with pathogen signalling. Some beneficial microbes may enter the root and disseminate in the aerial parts, such as the bacterial endophyte *Burkholderia phytofirmans*. The circulating microorganisms, that may also be pathogens like the bacteria *Xylella fastidiosa*, can reach leaves and integrate their highly dynamic microbiome, driven by seasonal variations. Throughout the vegetative season, leaves are colonised by major grapevine pathogens, such as the causal agents of mildews (*Plasmopara viticola* and *Erysiphe necator*) which are both introduced pathogens of North-American origin. Leaves also harbour microorganisms with biocontrol activity (e.g., *Bacillus subtilis*, *Aureobasidium pullulans*, *Ampelomyces* spp.). In mature grapevines, wood degradation associated with complex sets of microorganisms (e.g., *Phaeoconiella* spp., *Fomitiporia* spp., *Eutypa lata*, *Botryosphaeriaceae* spp.) may occur and lead to plant death. The search for biocontrol agents among mycoviruses, bacteria and fungi present in the wood is an active field of research requiring the inference of microbial interaction networks. Finally, the berry microbiome varies markedly during the development of the berry and contains, at the ripening stage, some fermentative yeasts and bacteria (e.g., *Saccharomyces cerevisiae*, *Oenococcus oeni*) positively involved in winemaking, but also undesirable microorganisms such as fungi producing off-flavours (e.g., the fungal pathogen *Botrytis cinerea*) and acetic acid bacteria leading to wine defaults.

Omics and associated computational approaches can further improve our understanding of the functioning of the grapevine microbiome, to ultimately design beneficial microbial consortia that could be inoculated to foster grapevine performance (health, growth and berry quality) under conditions of climate change and reduced pesticide use. To reach this objective, we propose to identify beneficial microbial taxa by performing field sampling campaigns informed by grapevine evolutionary history and phenotypic measurements (disease resistance, drought tolerance). Statistical approaches could then be applied to the metabarcoding datasets to identify subsets of microbial taxa whose presence or abundance is significantly related to grapevine phenotypic variations. Shotgun metagenomics could then be used to uncover microbial genomes and functions in selected samples, while microfluidic qPCR chips could be used to accurately estimate the abundance of selected microbial taxa. These data would then feed mathematical models to decipher metabolic and ecological interactions in small

microbial consortia and analyse how these interactions drive microbial community dynamics and variations in plant phenotypic traits. The high-throughput cultivation of environmental samples in parallel with sequencing will be crucial to isolate microbial strains of interest and experimentally validate hypotheses generated by mathematical models.

Based on current knowledge and available methods, we propose the following research avenues to make the grapevine microbiome more functional:

**A.** *Design microbiome studies informed by evolutionary history*

- Analyse the microbiome of wild grapevine and ancestral grapevine varieties to discover beneficial microorganisms that may have been lost during domestication and breeding
- Analyse the microbiome of American *Vitis* species to discover potential antagonists of North-American pathogens

**B.** *Consider grapevine as a dynamic system of interactions*

- Investigate how rootstock-scion interactions in the graft zone shape the microbiome of the whole plant
- Decipher the microbial interactions that regulate the dynamics of pathogen populations
- Assess the cascading effects of the soil microbiome on the root, trunk, leaf and berry microbiome, and their consequences on plant health
- Analyse the microbial fluxes that govern the seasonal dynamics of the berry microbiome, and their consequences on wine properties
- Identify and monitor the microbiome changes that precede the development of trunk diseases
- Identify the viruses (mycoviruses and phages, in particular) that shape the dynamics of the grapevine microbiome

**C.** *Combine metagenomics, culturomics and computational approaches*

- Use field experiments and metabarcoding data to highlight small consortia of microbial taxa related to variations in grapevine phenotypic traits
- Create and maintain microbial culture collections to perform controlled experiments with the selected microbial consortia
- Develop shotgun metagenomics and metabolic network modelling to understand the functioning of those microbial consortia
- Monitor selected microbial populations with quantitative approaches to model and predict their dynamics in response to environmental changes and management scenarios.

These research avenues are intertwined and not independent of each other: they require the creation of multidisciplinary research consortia with complementary expertise to define the scientific questions, experimental designs and analysis pipelines.

## Acknowledgements

We are very grateful to Maureen Pellan for the design of Fig. 3. The authors acknowledge the support of the French National Research Agency (ANR) under the grants 20-PCPA-0010 (PPR VITAE) and ANR-17-CE32-0011 (NGB), and of FranceAgrimer and CNIV under the grant n°2018-52537 (Vitirhizobiome). LP received funding from *Fondation Bordeaux Université*.

## References

- Abdelfattah, A., Sanzani, S.M., Wisniewski, M., Berg, G., Cacciola, S.O., Schena, L., 2019. Revealing cues for fungal interplay in the plant–air interface in vineyards. *Front. Plant Sci.* 10, 922.
- Abrego, N., Roslin, T., Huotari, T., Tack, A.J.M., Lindahl, B.D., Tikhonov, G., Somervuo, P., Schmidt, N.M., Ovaskainen, O., 2020. Accounting for environmental variation in co-occurrence modelling reveals the importance of positive interactions in root-associated fungal communities. *Mol. Ecol.* 29, 2736–2746.
- Aleynova, O.A., Nityagovsky, N.N., Dubrovina, A.S., Kiselev, K.V., 2022. The biodiversity of grapevine bacterial endophytes of *Vitis amurensis* Rupr. *Plan. Theory* 11, 1128.
- Alkowni, R., Zhang, Y.-P., Rowhani, A., Uyemoto, J.K., Minafra, A., 2011. Biological, molecular, and serological studies of a novel strain of grapevine leafroll-associated virus 2. *Virus Genes* 43, 102–110.
- Anderson, K., Aryal, N.R., 2013. Which Winegrape Varieties are Grown Where? A Global Empirical Picture. University of Adelaide Press.
- Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S., Dehal, P., Ware, D., Perez, F., Canon, S., Sneddon, M.W., Henderson, M.L., Riehl, W.J., Murphy-Olson, D., Chan, S.Y., Kamimura, R.T., Kumari, S., Drake, M.M., Brettin, T.S., Glass, E.M., Chivian, D., Gunter, D., Weston, D.J., Allen, B.H., Baumohl, J., Best, A.A., Bowen, B., Brenner, S.E., Bun, C.C., Chandonia, J.-M., Chia, J.-M., Colasanti, R., Conrad, N., Davis, J.J., Davison, B.H., DeJongh, M., Devoid, S., Dietrich, E., Dubchak, I., Edirisinghe, J.N., Fang, G., Faria, J.P., Frybarger, P.M., Gerlach, W., Gerstein, M., Greiner, A., Gurtowski, J., Haun, H.L., He, F., Jain, R., Joachimiak, M.P., Keegan, K.P., Kondo, S., Kumar, V., Land, M.L., Meyer, F., Mills, M., Novichkov, P.S., Oh, T., Olsen, G.J., Olson, R., Parrello, B., Pasternak, S., Pearson, E., Poon, S.S., Price, G.A., Ramakrishnan, S., Ranjan, P., Ronald, P.C., Schatz, M.C., Seaver, S.M.D., Shukla, M., Sutormin, R.A., Syed, M.H., Thomason, J., Tintle, N.L., Wang, D., Xia, F., Yoo, H., Yoo, S., Yu, D., 2018. KBase: The United States department of energy systems biology knowledgebase. *Nat. Biotechnol.* 36, 566–569.
- Armijo, G., Schlechter, R., Agurto, M., Muñoz, D., Nuñez, C., Arce-Johnson, P., 2016. Grapevine pathogenic microorganisms: understanding infection strategies and host response scenarios. *Front. Plant Sci.* 7, 382.
- Armitage, D.W., Jones, S.E., 2019. How sample heterogeneity can obscure the signal of microbial interactions. *ISME J.* 13, 2639–2646.

- Arnold, C., Schnitzler, A., Douard, A., Peter, R., Gillet, F., 2005. Is there a future for wild grapevine (*Vitis vinifera* subsp. *silvestris*) in the Rhine Valley? *Biodivers. Conserv.* 14, 1507–1523.
- Arroyo-García, R., Ruiz-García, L., Bolling, L., Ocete, R., López, M.A., Arnold, C., Ergul, A., SöylemezoLu, G., Uzun, H.I., Cabello, F., Ibáñez, J., Aradhya, M.K., Atanassov, A., Atanassov, I., Balint, S., Cenis, J.L., Costantini, L., Gorislavets, S., Grando, M.S., Klein, B.Y., McGovern, P.E., Merdinoglu, D., Pejic, I., Pelsy, F., Primikiri, N., Risovannaya, V., Roubelakis-Angelakis, K.A., Snoussi, H., Sotiri, P., Tamhankar, S., This, P., Troshin, L., Malpica, J.M., Lefort, F., Martínez-Zapater, J.M., 2006. Multiple origins of cultivated grapevine (*Vitis vinifera* L. ssp. *sativa*) based on chloroplast DNA polymorphisms. *Mol. Ecol.* 15, 3707–3714.
- Asad, N.I., Tremblay, J., Dozois, J., Mukula, E., L'Espérance, E., Constant, P., Yergeau, E., 2022. Predictive microbial-based modelling of wheat yields and grain baking quality across a 500 km transect in Québec. *FEMS Microbiol. Ecol.* 97, fiab160.
- Asghari, S., Harighi, B., Mozafari, A.A., Esmael, Q., Ait Barka, E., 2019. Screening of endophytic bacteria isolated from domesticated and wild growing grapevines as potential biological control agents against crown gall disease. *BioControl* 64, 723–735.
- Asghari, S., Harighi, B., Ashengroph, M., Clement, C., Aziz, A., Esmael, Q., Ait Barka, E., 2020. Induction of systemic resistance to *Agrobacterium tumefaciens* by endophytic bacteria in grapevine. *Plant Pathol.* 69, 827–837.
- Azevedo-Silva, D., Rasmussen, J.A., Carneiro, M., Gilbert, M.T.P., Azevedo, H., 2021. Feasibility of applying shotgun metagenomic analyses to grapevine leaf, rhizosphere and soil microbiome characterisation. *Aust. J. Grape Wine Res.* 27, 519–526.
- Aziz, A., Verhagen, B., Magnin-Robert, M., Couderchet, M., Clément, C., Jeandet, P., Trotel-Aziz, P., 2016. Effectiveness of beneficial bacteria to promote systemic resistance of grapevine to gray mold as related to phytoalexin production in vineyards. *Plant and Soil* 405, 141–153.
- Aziz, A., Perazzolli, M., Gramaje, D., Zyprian, E.M., Cantu, D., 2020. Editorial: recent advances on grapevine-microbe interactions: from signal perception to resistance response. *Front. Plant Sci.* 11, 1164.
- Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S., Vivanco, J.M., 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* 57, 233–266.
- Baker, M.E., King, R.S., Kahle, D., 2015. TITAN2: Threshold Indicator Taxa Analysis. R Package Version 2.1.
- Bakshi, S., Szejnberg, A., Yarden, O., 2001. Isolation and characterization of a cold-tolerant strain of *Fusarium proliferatum*, a biocontrol agent of grape downy mildew. *Phytopathology* 91, 1062–1068.
- Baldan, E., Nigris, S., Romualdi, C., D'Alessandro, S., Clocchiatti, A., Zottini, M., Stevanato, P., Squartini, A., Baldan, B., 2015. Beneficial bacteria isolated from grapevine inner tissues shape *Arabidopsis thaliana* roots. *Plos one* 10, e0140252.
- Balestrini, R., Magurno, F., Walker, C., Lumini, E., Bianciotto, V., 2010. Cohorts of arbuscular mycorrhizal fungi (AMF) in *Vitis vinifera*, a typical Mediterranean fruit crop: cohorts of AM fungi in vineyards. *Environ. Microbiol. Rep.* 2, 594–604.
- Bao, L., Sun, B., Wei, Y., Xu, N., Zhang, S., Gu, L., Bai, Z., 2022. Grape cultivar features differentiate the grape rhizosphere microbiota. *Plants* 11, 1111.
- Barata, A., Malfeito-Ferreira, M., Loureiro, V., 2012. The microbial ecology of wine grape berries. *Int. J. Food Microbiol.* 153, 243–259.
- Barroso-Bergadà, D., Pauvert, C., Vallance, J., Delière, L., Bohan, D.A., Buée, M., Vacher, C., 2021. Microbial networks inferred from environmental DNA data for biomonitoring ecosystem change: Strengths and pitfalls. *Mol. Ecol. Resour.* 21, 762–780.

- Barroso-Bergada, D., Tamaddoni-Nezhad, A., Muggleton, S.H., Vacher, C., Galic, N., Bohan, D.A., 2022. Machine learning of microbial interactions using abductive ILP and hypothesis frequency/compression estimation. In: Katzouris, N., Artikis, A. (Eds.), *Inductive Logic Programming. Lecture Notes in Computer Science*, Springer International Publishing, Cham, pp. 26–40.
- Becerra-Lucio, A.A., Labrín-Sotomayor, N.Y., Becerra-Lucio, P.A., Trujillo-Elisea, F.I., Chávez-Bárceñas, A.T., Machkour-M'Rabet, S., Peña-Ramírez, Y.J., 2021. Diversity and interactomics of bacterial communities associated with dominant trees during tropical forest recovery. *Curr. Microbiol.* 78, 3417–3429.
- Beghini, F., McIver, L.J., Blanco-Míguez, A., Dubois, L., Asnicar, F., Maharjan, S., Mailyan, A., Manghi, P., Scholz, M., Thomas, A.M., Valles-Colomer, M., Weingart, G., Zhang, Y., Zolfo, M., Huttenhower, C., Franzosa, E.A., Segata, N., 2021. Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife* 10, e65088.
- Behrens, F.H., Fischer, M., 2022. Evaluation of different phyllosphere sample types for parallel metabarcoding of fungi and oomycetes in *Vitis vinifera*. *Phytobiomes J.* 6, 207–213 (PBIOMES-11-21-0072-SC).
- Bekris, F., Vasileiadis, S., Papadopoulou, E., Samaras, A., Testempasis, S., Gkizi, D., Tavlaki, G., Tzima, A., Paplomatas, E., Markakis, E., Karaoglanidis, G., Papadopoulou, K.K., Karpouzias, D.G., 2021. Grapevine wood microbiome analysis identifies key fungal pathogens and potential interactions with the bacterial community implicated in grapevine trunk disease appearance. *Environ. Microbiome* 16, 23.
- Belcour, A., Frioux, C., Aite, M., Bretaudeau, A., Hildebrand, F., Siegel, A., 2020. Metage2Metabo, microbiota-scale metabolic complementarity for the identification of key species. *Elife* 9, e61968.
- Belcour, A., Ruiz, B., Frioux, C., Blanquart, S., Siegel, A., 2022. EsMcCaTa: Estimating metabolic capabilities from taxonomic affiliations. *bioRxiv*. 2022.03.16.484574.
- Belessi, C., Chalvanti, I., Marmaras, I., Nisiotou, A., 2022. The effect of vine variety and vintage on wine yeast community structure of grapes and ferments. *J. Appl. Microbiol.* 132, 3672–3684.
- Berg, G., Rybakova, D., Fischer, D., Cernava, T., Champomier Vergès, M.-C., Charles, T., Chen, X., Cocolin, L., Eversole, K., Corral, G.H., Kazou, M., Kinkel, L., Lange, L., Lima, N., Loy, A., Macklin, J.A., Maguin, E., Mauchline, T., McClure, R., Mitter, B., Ryan, M., Sarand, I., Smidt, H., Schelkle, B., Roume, H., Kiran, G.S., Selvin, J., de Souza, R.S.C., van Overbeek, L., Singh, B.K., Wagner, M., Walsh, A., Sessitsch, A., Schloter, M., 2020. Microbiome definition re-visited: old concepts and new challenges. *Microbiome* 8, 103.
- Berlanas, C., Berbegal, M., Elena, G., Laidani, M., Cibriain, J.F., Sagües, A., Gramaje, D., 2019. The fungal and bacterial rhizosphere microbiome associated with grapevine rootstock genotypes in mature and young vineyards. *Front. Microbiol.* 10, 1142.
- Bertsch, C., Ramírez-Suero, M., Magnin-Robert, M., Larignon, P., Chong, J., Abou-Mansour, E., Spagnolo, A., Clément, C., Fontaine, F., 2013. Grapevine trunk diseases: complex and still poorly understood: *Grapevine trunk diseases*. *Plant Pathol.* 62, 243–265.
- Bettenfeld, P., Cadenai Canals, J., Jacquens, L., Fernandez, O., Fontaine, F., van Schaik, E., Courty, P.-E., Trouvelot, S., 2021. The microbiota of the grapevine holobiont: a key component of plant health. *J. Adv. Res.* 40, 1–15. S2090123221002617.
- Bhowmick, T.S., Das, M., Heinz, K.M., Krauter, P.C., Gonzalez, C.F., 2016. Transmission of phage by glassy-winged sharpshooters, a vector of *Xylella fastidiosa*. *Bacteriophage* 6, e1218411.



- Bickhart, D.M., Kolmogorov, M., Tseng, E., Portik, D.M., Korobeynikov, A., Tolstoganov, I., Uritskiy, G., Liachko, I., Sullivan, S.T., Shin, S.B., Zorea, A., Andreu, V.P., Panke-Buisse, K., Medema, M.H., Mizrahi, I., Pevzner, P.A., Smith, T.P.L., 2022. Generating lineage-resolved, complete metagenome-assembled genomes from complex microbial communities. *Nat. Biotechnol.* 40, 1–9.
- Biget, M., 2021. La vigne comme holobionte: validité du concept, dynamique temporelle, déterminants environnementaux et application à la définition du terroir microbiologique. Doctoral dissertation. University of Rennes 1, 74–104.
- Biggs, M.B., Medlock, G.L., Kolling, G.L., Papin, J.A., 2015. Metabolic network modeling of microbial communities. *Wiley Interdiscip. Rev. Syst. Biol. Med.* 7, 317–334.
- Bilen, M., Dufour, J.-C., Lagier, J.-C., Cadoret, F., Daoud, Z., Dubourg, G., Raoult, D., 2018. The contribution of culturomics to the repertoire of isolated human bacterial and archaeal species. *Microbiome* 6, 94.
- Bleach, C., Ridgway, H.J., Jaspers, M.V., Jones, E.E., 2021. The relative susceptibility of grapevine rootstocks to black foot disease is dependent on inoculum pressure. *N. Z. Plant Prot.* 74, 62–69.
- Bodin, E., Bellée, A., Dufour, M.-C., André, O., Corio-Costet, M.-F., 2020. Grapevine stimulation: a multidisciplinary approach to investigate the effects of biostimulants and a plant defense stimulator. *J. Agric. Food Chem.* 68, 15085–15096.
- Bois, B., Zito, S., Calonne, A., 2017. Climate vs grapevine pests and diseases worldwide: the first results of a global survey. *OENO One* 51, 133–139.
- Bokulich, N.A., Thorngate, J.H., Richardson, P.M., Mills, D.A., 2014. Microbial biogeography of wine grapes is conditioned by cultivar, vintage, and climate. *Proc. Natl. Acad. Sci. U. S. A.* 111, 139–148.
- Bona, E., Massa, N., Novello, G., Boatti, L., Cesaro, P., Todeschini, V., Magnelli, V., Manfredi, M., Marengo, E., Mignone, F., Berta, G., Lingua, G., Gamalero, E., 2018. Metaproteomic characterization of *Vitis vinifera* rhizosphere. *FEMS Microbiol. Ecol.* 95, 1–15.
- Borcard, D., Legendre, P., 2002. All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecol. Model.* 153, 51–68.
- Borderes, M., Gasc, C., Prestat, E., Galvão Ferrarini, M., Vinga, S., Boucinha, L., Sagot, M.-F., 2021. A comprehensive evaluation of binning methods to recover human gut microbial species from a non-redundant reference gene catalog. *NAR Genom. Bioinform.* 3, lqab009.
- Börlin, M., Claisse, O., Albertin, W., Salin, F., Legras, J.L., Masneuf-Pomaredé, I., 2020. Quantifying the effect of human practices on *S. cerevisiae* vineyard metapopulation diversity. *Sci. Rep.* 10 (1), 16214.
- Botstein, D., Fink, G.R., 2011. yeast: an experimental organism for 21st century biology. *Genetics* 189, 695–704.
- Bowers, J., Boursiquot, J.-M., This, P., Chu, K., Johansson, H., Meredith, C., 1999. Historical genetics: the parentage of chardonnay, gamay, and other wine grapes of Northeastern France. *Science* 285, 1562–1565.
- Bozoudi, D., Tsaltas, D., 2018. The multiple and versatile roles of *Aureobasidium pullulans* in the vitivicultural sector. *Fermentation* 4, 85.
- Brewer, M.T., Milgroom, M.G., 2010. Phylogeography and population structure of the grape powdery mildew fungus, *Erysiphe necator*, from diverse *Vitis* species. *BMC Evol. Biol.* 10, 268.
- Bruetz, E., Lecomte, P., Grosman, J., Doublet, B., Bertsch, C., Ugaglia, A., Teissedre, P.-L., Costa, J.-P.D., Guerin-Dubrana, L., Rey, P., 2013. Overview of grapevine trunk diseases in France in the 2000s. *Phytopathol. Mediterr.* 52, 262–275.

- Bruetz, E., Vallance, J., Gerbore, J., Lecomte, P., Da Costa, J.-P., Guerin-Dubrana, L., Rey, P., 2014. Analyses of the temporal dynamics of fungal communities colonizing the healthy wood tissues of esca leaf-symptomatic and asymptomatic vines. *PLoS One* 9, e95928.
- Bruetz, E., Haidar, R., Alou, M.T., Vallance, J., Bertsch, C., Mazet, F., Fermaud, M., Deschamps, A., Guerin-Dubrana, L., Compant, S., Rey, P., 2015. Bacteria in a wood fungal disease: characterization of bacterial communities in wood tissues of esca-foliar symptomatic and asymptomatic grapevines. *Front. Microbiol.* 6, 1137.
- Bruetz, E., Baumgartner, K., Bastien, S., Travadon, R., Guérin-Dubrana, L., Rey, P., 2016. Various fungal communities colonise the functional wood tissues of old grapevines externally free from grapevine trunk disease symptoms: fungal microflora of GTD-free old vines. *Aust. J. Grape Wine Res.* 22, 288–295.
- Bruetz, E., Vallance, J., Gautier, A., Laval, V., Compant, S., Maurer, W., Sessitsch, A., Lebrun, M., Rey, P., 2020. Major changes in grapevine wood microbiota are associated with the onset of esca, a devastating trunk disease. *Environ. Microbiol.* 22, 5189–5206.
- Bruisson, S., Zufferey, M., L'Haridon, F., Trutmann, E., Anand, A., Dutartre, A., De Vrieze, M., Weisskopf, L., 2019. Endophytes and epiphytes from the grapevine leaf microbiome as potential biocontrol agents against phytopathogens. *Front. Microbiol.* 10, 2726.
- Brum, M.C.P., Araújo, W.L., Maki, C.S., Azevedo, J.L., 2012. Endophytic fungi from *Vitis labrusca* L. ('Niagara Rosada') and its potential for the biological control of *Fusarium oxysporum*. *Genet. Mol. Res.* 11, 4187–4197.
- Bucci, V., Tzen, B., Li, N., Simmons, M., Tanoue, T., Bogart, E., Deng, L., Yeliseyev, V., Delaney, M.L., Liu, Q., Olle, B., Stein, R.R., Honda, K., Bry, L., Gerber, G.K., 2016. MDSINE: microbial dynamical systems INference engine for microbiome time-series analyses. *Genome Biol.* 17, 121.
- Burruano, S., 2000. The life-cycle of *Plasmopara viticola*, cause of downy mildew of vine. *Mycologist* 14, 179–182.
- Burruano, S., Alfonso, A., Lo Piccolo, S., Conigliaro, G., Mondello, V., Torta, L., Moretti, M., Assante, G., 2008. Interaction between *Acremonium byssoides* and *Plasmopara viticola* in *Vitis vinifera*. *Phytopathol. Mediterr.* 47, 122–131.
- Burruano, S., Mondello, V., Conigliaro, G., 2016. Endophytic fungi in asymptomatic *Vitis vinifera* L. and their effects on *Plasmopara viticola*. In: Compant, S., Mathieu, F. (Eds.), *Biocontrol of Major Grapevine Diseases: Leading Research*. CABI, Wallingford, pp. 98–112.
- Busby, P.E., Soman, C., Wagner, M.R., Friesen, M.L., Kremer, J., Bennett, A., Morsy, M., Eisen, J.A., Leach, J.E., Dangl, J.L., 2017. Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol.* 15, e2001793.
- Button, D.K., Schut, F., Quang, P., Martin, R., Robertson, B.R., 1993. Viability and isolation of marine bacteria by dilution culture: theory, procedures, and initial results. *Appl. Environ. Microbiol.* 59, 881–891.
- Cadle-Davidson, L., Chicoine, D.R., Consolie, N.H., 2011. Variation within and among *Vitis* spp. for foliar resistance to the powdery mildew pathogen *Erysiphe necator*. *Plant Dis.* 95, 202–211.
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13, 581–583.
- Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* 11, 2639–2643.
- Calonne, A., Cartolaro, P., Naulin, J.-M., Bailey, D., Langlais, M., 2008. A host-pathogen simulation model: powdery mildew of grapevine. *Plant Pathol.* 57, 493–508.

- Calonnet, A., Jolivet, J., Vivin, P., Schnee, S., 2018. Pathogenicity traits correlate with the susceptible *Vitis vinifera* leaf physiology transition in the biotroph fungus *Erysiphe necator*: an adaptation to plant ontogenic resistance. *Front. Plant Sci.* 9, 1808.
- Cambon, M.C., Cartry, D., Chancerel, E., Ziegler, C., Levionnois, S., Coste, S., Stahl, C., Delzon, S., Buée, M., Burban, B., Casal, J., Fort, T., Goret, J.-Y., Heuret, P., Léger, P., Louisanna, E., Ritter, Y., Bonal, D., Roy, M., Schimann, H., Vacher, C., 2022. Drought tolerance traits in neotropical trees correlate with the composition of phyllosphere fungal communities. *Phytobiomes J. (PBIOMES-04-22-0023-R)*.
- Campisano, A., Antonielli, L., Pancher, M., Yousaf, S., Pindo, M., Pertot, I., 2014. Bacterial endophytic communities in the grapevine depend on pest management. *PLoS One* 9, e112763.
- Campisano, A., Pancher, M., Puopolo, G., Puddu, A., López-Fernández, S., Biagini, B., Yousaf, S., Pertot, I., 2015. Diversity in endophyte populations reveals functional and taxonomic diversity between wild and domesticated grapevines. *Am. J. Enol. Vitic.* 66, 12–21.
- Canfora, L., Vendramin, E., Felici, B., Tarricone, L., Florio, A., Benedetti, A., 2018. Vineyard microbiome variations during different fertilisation practices revealed by 16s rRNA gene sequencing. *Appl. Soil Ecol.* 125, 71–80.
- Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol. Biol. Evol.* 38, msab293.
- Carbone, M.J., Alaniz, S., Mondino, P., Gelabert, M., Eichmeier, A., Tekielska, D., Bujanda, R., Gramaje, D., 2021. Drought influences fungal community dynamics in the grapevine rhizosphere and root microbiome. *J. Fungi* 7, 686.
- Carmichael, P.C., Siyoum, N., Chidamba, L., Korsten, L., 2019. Exploring the microbial communities associated with *Botrytis cinerea* during berry development in table grape with emphasis on potential biocontrol yeasts. *Eur. J. Plant Pathol.* 154, 919–930.
- Carr, A., Diener, C., Baliga, N.S., Gibbons, S.M., 2019. Use and abuse of correlation analyses in microbial ecology. *ISME J.* 13, 2647–2655.
- Carrere, C., Spilmont, A.-S., Loupuit, G., Beutin, P., Stessels, C., Ollat, N., Cookson, S.-J., 2022. Evaluation of criteria to assist the selection of good quality grafted grapevines prior to their commercialisation. *OENO One* 56, 15–27.
- Castañeda, L.E., Barbosa, O., 2017. Metagenomic analysis exploring taxonomic and functional diversity of soil microbial communities in Chilean vineyards and surrounding native forests. *PeerJ* 5, e3098.
- Castañeda, L.E., Miura, T., Sánchez, R., Barbosa, O., 2018. Effects of agricultural management on phyllosphere fungal diversity in vineyards and the association with adjacent native forests. *PeerJ* 6, e5715.
- Cavaliere, D., McGovern, P.E., Hartl, D.L., Mortimer, R., Polsinelli, M., 2003. Evidence for *S. cerevisiae* fermentation in ancient wine. *J. Mol. Evol.* 57, S226–S232.
- Chandra, M., Mota, M., Silva, A.C., Malfeito-Ferreira, M., 2020. Forest oak woodlands and fruit tree soils are reservoirs of wine-related yeast species. *Am. J. Enol. Vitic.* 71, 191–197.
- Chaparro, J.M., Badri, D.V., Vivanco, J.M., 2014. Rhizosphere microbiome assemblage is affected by plant development. *ISME J.* 8, 790–803.
- Chaudhry, V., Runge, P., Sengupta, P., Doehlemann, G., Parker, J.E., Kemen, E., 2021. Shaping the leaf microbiota: plant–microbe–microbe interactions. *J. Exp. Bot.* 72, 36–56.
- Chen, F., Guo, Y.B., Wang, J.H., Li, J.Y., Wang, H.M., 2007. Biological control of grape crown gall by *Rahnella aquatilis* HX2. *Plant Dis.* 91, 957–963.
- Chen, M., Brun, F., Raynal, M., Makowski, D., 2019. Timing of grape downy mildew onset in bordeaux vineyards. *Phytopathology* 109, 787–795.

- Chen, L.-X., Anantharaman, K., Shaiber, A., Eren, A.M., Banfield, J.F., 2020. Accurate and complete genomes from metagenomes. *Genome Res.* 30, 315–333.
- Chen, Y.-H., Chiang, P.-W., Rogozin, D.Y., Degermendzhy, A.G., Chiu, H.-H., Tang, S.-L., 2021. Salvaging high-quality genomes of microbial species from a meromictic lake using a hybrid sequencing approach. *Commun. Biol.* 4, 996.
- Chiapello, M., Rodríguez-Romero, J., Ayllón, M.A., Turina, M., 2020. Analysis of the virome associated to grapevine downy mildew lesions reveals new mycovirus lineages. *Virus Evol.* 6, veaa058.
- Chiquet, J., Robin, S., Mariadassou, M., 2019. Variational inference for sparse network reconstruction from count data. In: Proceedings of the 36th International Conference on Machine Learning. Presented at the International Conference on Machine Learning. PMLR, pp. 1162–1171.
- Chitwood, D.H., Klein, L.L., O’Hanlon, R., Chacko, S., Greg, M., Kitchen, C., Miller, A.J., Londo, J.P., 2016. Latent developmental and evolutionary shapes embedded within the grapevine leaf. *New Phytol.* 210, 343–355.
- Claisse, O., Chaïb, A., Jaomanjaka, F., Philippe, C., Barchi, Y., Lucas, P.M., Le Marrec, C., 2021. Distribution of prophages in the *Oenococcus oeni* species. *Microorganisms* 9, 856.
- Clavijo-Coppens, F., Ginet, N., Cesbron, S., Briand, M., Jacques, M.-A., Ansaldo, M., 2021. Novel virulent bacteriophages infecting mediterranean isolates of the plant pest *Xylella fastidiosa* and *Xanthomonas albilineans*. *Viruses* 13, 725.
- Cobo-Díaz, J.F., Legrand, F., Le Floch, G., Picot, A., 2022. Influence of maize residues in shaping soil microbiota and *Fusarium* spp. communities. *Microb. Ecol.* 83, 702–713.
- Cobos, R., Ibañez, A., Díez-Galán, A., Calvo-Peña, C., Ghoshshizadeh, S., Coque, J.J.R., 2022. The grapevine microbiome to the rescue: implications for the biocontrol of trunk diseases. *Plan. Theory* 11, 840.
- Compant, S., Kaplan, H., Sessitsch, A., Nowak, J., Ait Barka, E., Clément, C., 2008. Endophytic colonization of *Vitis vinifera* L. by *Burkholderia phytofirmans* strain PsJN: from the rhizosphere to inflorescence tissues. *FEMS Microbiol. Ecol.* 63, 84–93.
- Compant, S., Brader, G., Muzammil, S., Sessitsch, A., Lebrühi, A., Mathieu, F., 2013. Use of beneficial bacteria and their secondary metabolites to control grapevine pathogen diseases. *BioControl* 58, 435–455.
- Compant, S., Samad, A., Faist, H., Sessitsch, A., 2019. A review on the plant microbiome: ecology, functions, and emerging trends in microbial application. *J. Adv. Res.* 19, 29–37.
- Cookson, S.J., Clemente Moreno, M.J., Hevin, C., Nyamba Mendome, L.Z., Delrot, S., Trossat-Magnin, C., Ollat, N., 2013. Graft union formation in grapevine induces transcriptional changes related to cell wall modification, wounding, hormone signalling, and secondary metabolism. *J. Exp. Bot.* 64, 2997–3008.
- Cordero-Bueso, G., Arroyo, T., Serrano, A., Tello, J., Aporta, I., Velez, M., Valero, E., 2011a. Influence of the farming system and vine variety on yeast communities associated with grape berries. *Int. J. Food Microbiol.* 145, 132–139.
- Cordero-Bueso, G., Arroyo, T., Serrano, A., Valero, E., 2011b. Influence of different floor management strategies of the vineyard on the natural yeast population associated with grape berries. *Int. J. Food Microbiol.* 148, 23–29.
- Cordero-Bueso, G., Arroyo, T., Valero, E., 2014. A long term field study of the effect of fungicides penconazole and sulfur on yeasts in the vineyard. *Int. J. Food Microbiol.* 189, 189–194.
- Cordero-Bueso, G., Mangieri, N., Maghradze, D., Foschino, R., Valdetara, F., Cantoral, J.M., Vigentini, I., 2017. Wild grape-associated yeasts as promising biocontrol agents against *Vitis vinifera* fungal pathogens. *Front. Microbiol.* 8, 2025.
- Cordero-Bueso, G., Vigentini, I., Foschino, R., Maghradze, D., Ruiz-Muñoz, M., Benitez-Trujillo, F., Cantoral, J.M., 2022. Culturable yeast diversity of grape berries from *Vitis vinifera* ssp. *sylvestris* (Gmelin) hegi. *J. Fungi* 8, 410.

- Crane, S.L., van Dorst, J., Hose, G.C., King, C.K., Ferrari, B.C., 2018. Microfluidic qPCR enables high throughput quantification of microbial functional genes but requires strict curation of primers. *Front. Environ. Sci.* 6, 145.
- Crisp, P., Wicks, T.J., Bruer, D., Scott, E.S., 2006. An evaluation of biological and abiotic controls for grapevine powdery mildew. 2. Vineyard trials. *Aust. J. Grape Wine Res.* 12, 203–211.
- Cruz-Silva, A., Figueiredo, A., Sebastiana, M., 2021. First insights into the effect of mycorrhizae on the expression of pathogen effectors during the infection of grapevine with *Plasmopara viticola*. *Sustainability* 13, 1226.
- Cureau, N., Threlfall, R., Marasini, D., Lavefve, L., Carbonero, F., 2021. Year, location, and variety impact on grape-associated mycobiota of arkansas-grown wine grapes for wine production. *Microb. Ecol.* 82, 845–858.
- Dagostin, S., Schärer, H.-J., Pertot, I., Tamm, L., 2011. Are there alternatives to copper for controlling grapevine downy mildew in organic viticulture? *Crop Prot.* 30, 776–788.
- Damme, R.V., Hölzer, M., Viehweger, A., Müller, B., Bongcam-Rudloff, E., Brandt, C., 2021. Metagenomics workflow for hybrid assembly, differential coverage binning, metatranscriptomics and pathway analysis (MUFFIN). *PLoS Comput. Biol.* 17, e1008716.
- Darriaut, R., Martins, G., Dewasme, C., Mary, S., Darrieutort, G., Ballestra, P., Marguerit, E., Vivin, P., Ollat, N., Masneuf-Pomarède, I., Lauvergeat, V., 2021. Grapevine decline is associated with difference in soil microbial composition and activity. *OENO One* 55, 67–84.
- Darriaut, R., Lailheugue, V., Masneuf-Pomarède, I., Marguerit, E., Martins, G., Compant, S., Ballestra, P., Upton, S., Ollat, N., Lauvergeat, V., 2022. Grapevine rootstock and soil microbiome interactions: keys for a resilient viticulture. *Hortic. Res.* 9, uhac019.
- Das, M., Bhowmick, T.S., Ahern, S.J., Young, R., Gonzalez, C.F., 2015. Control of pierce's disease by phage. *PLoS One* 10, e0128902.
- de Brito, L.M., Zanette, F., Biasi, L.A., Costa, M.D., dos Passos, J.F.M., 2019. Morphological identification of *Ilyonectria liriodendri* and its interaction with plant growth-promoting bacteria in grapevine rootstocks. *Rev. Bras. Frutic.* 41, e-140.
- Del Frari, G., Gobbi, A., Aggerbeck, M.R., Oliveira, H., Hansen, L.H., Ferreira, R.B., 2019. Characterization of the wood mycobiome of *vitis vinifera* in a vineyard affected by esca. Spatial distribution of fungal communities and their putative relation with leaf symptoms. *Front. Plant Sci.* 10, 910.
- del Martínez-Diz, M.P., Andrés-Sodupe, M., Bujanda, R., Díaz-Losada, E., Eichmeier, A., Gramaje, D., 2019. Soil-plant compartments affect fungal microbiome diversity and composition in grapevine. *Fungal Ecol.* 41, 234–244.
- Derocles, S.A.P., Bohan, D.A., Dumbrell, A.J., Kitson, J.J.N., Massol, F., Pauvert, C., Plantegenest, M., Vacher, C., Evans, D.M., 2018. Biomonitoring for the 21st century: integrating next-generation sequencing into ecological network analysis. In: *Advances in Ecological Research*. Elsevier, pp. 1–62.
- Deyett, E., Rolshausen, P.E., 2019. Temporal dynamics of the sap microbiome of grapevine under high pierce's disease pressure. *Front. Plant Sci.* 10, 1246.
- Deyett, E., Rolshausen, P.E., 2020. Endophytic microbial assemblage in grapevine. *FEMS Microbiol. Ecol.* 96, faa053.
- Di Vecchi-Staraz, M., Laucou, V., Bruno, G., Lacombe, T., Gerber, S., Bourse, T., Boselli, M., This, P., 2009. Low level of pollen-mediated gene flow from cultivated to wild grapevine: consequences for the evolution of the endangered subspecies *Vitis vinifera* L. subs. *silvestris*. *J. Hered.* 100, 66–75.

- Ding, Y., Wei, R., Wang, L., Yang, C., Li, H., Wang, H., 2021. Diversity and dynamics of microbial ecosystem on berry surface during the ripening of Ecolly (*Vitis vinifera* L.) grape in Wuhai, China. *World J. Microbiol. Biotechnol.* 37, 214.
- Dohlman, A.B., Shen, X., 2019. Mapping the microbial interactome: statistical and experimental approaches for microbiome network inference. *Exp. Biol. Med.* 244, 445–458.
- Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M., Huttenhower, C., Langille, M.G.I., 2020. PICRUSt2 for prediction of metagenome functions. *Nat. Biotechnol.* 38, 685–688.
- Dreier, M., Berthoud, H., Shani, N., Wechsler, D., Junier, P., 2021. Development of a high-throughput microfluidic qPCR system for the quantitative determination of quality-relevant bacteria in cheese. *Front. Microbiol.* 11, 619166.
- Dries, L., Bussotti, S., Pozzi, C., Kunz, R., Schnell, S., Löhnertz, O., Vortkamp, A., 2021. Rootstocks shape their microbiome—bacterial communities in the rhizosphere of different grapevine rootstocks. *Microorganisms* 9, 822.
- Dufour, M.-C., Magnin, N., Dumas, B., Vergnes, S., Corio-Costet, M.-F., 2016. Highthroughput gene-expression quantification of grapevine defense responses in the field using microfluidic dynamic arrays. *BMC Genomics* 17, 957.
- Ebenhöh, O., Handorf, T., Heinrich, R., 2004. Structural analysis of expanding metabolic networks. *Genome Inform. Int. Conf. Genome Inform.* 15, 35–45.
- Eder, M., Nidelet, T., Sanchez, I., Camarasa, C., Legras, J.-L., Dequin, S., 2020. QTL mapping of modelled metabolic fluxes reveals gene variants impacting yeast central carbon metabolism. *Sci. Rep.* 10, 2162.
- Eichmeier, A., Pečenka, J., Peňázová, E., Baránek, M., Català-García, S., León, M., Armengol, J., Gramaje, D., 2018. High-throughput amplicon sequencing-based analysis of active fungal communities inhabiting grapevine after hot-water treatments reveals unexpectedly high fungal diversity. *Fungal Ecol.* 36, 26–38.
- Elena, G., Bruez, E., Rey, P., Luque, J., 2018. Microbiota of grapevine woody tissues with or without esca-foliar symptoms in northeast Spain. *Phytopathol. Mediterr.* 57, 425–438.
- Esmael, Q., Jacquard, C., Sanchez, L., Clément, C., Ait Barka, E., 2020. The mode of action of plant associated Burkholderia against grey mould disease in grapevine revealed through traits and genomic analyses. *Sci. Rep.* 10, 19393.
- Estaki, M., Jiang, L., Bokulich, N.A., McDonald, D., González, A., Kosciolk, T., Martino, C., Zhu, Q., Birmingham, A., Vázquez-Baeza, Y., Dillon, M.R., Bolyen, E., Caporaso, J.G., Knight, R., 2020. QIIME 2 enables comprehensive end-to-end analysis of diverse microbiome data and comparative studies with publicly available data. *Curr. Protoc. Bioinformatics* 70, 100.
- Falk, S.P., 1995. Parasitism of *Uncinula necator* cleistothecia by the mycoparasite *Ampelomyces quisqualis*. *Phytopathology* 85, 794.
- Falk, S., 1996. *Fusarium proliferatum* as a biocontrol agent against grape downy mildew. *Phytopathology* 86, 1010.
- Fan, Y., Gao, L., Chang, P., Li, Z., 2020. Endophytic fungal community in grape is correlated to foliar age and domestication. *Ann. Microbiol.* 70, 30.
- Fang, H., Huang, C., Zhao, H., Deng, M., 2015. CClasso: correlation inference for compositional data through Lasso. *Bioinformatics* 31, 3172–3180.
- Faust, K., Bauchinger, F., Laroche, B., de Buyl, S., Lahti, L., Washburne, A.D., Gonze, D., Widder, S., 2018. Signatures of ecological processes in microbial community time series. *Microbiome* 6, 120.
- Fay, J.C., Benavides, J.A., 2005. Evidence for domesticated and wild populations of *Saccharomyces cerevisiae*. *PLoS Genet.* 1, e5.
- Fay, J.C., McCullough, H.L., Sniegowski, P.D., Eisen, M.B., 2004. Population genetic variation in gene expression is associated with phenotypic variation in *Saccharomyces cerevisiae*. *Genome Biol.* 5, R26.

- Feechan, A., Kocsis, M., Riaz, S., Zhang, W., Gadoury, D.M., Walker, M.A., Dry, I.B., Reisch, B., Cadle-Davidson, L., 2015. Strategies for *RUN1* deployment using *RUN2* and *REN2* to manage grapevine powdery mildew informed by studies of race specificity. *Phytopathology* 105, 1104–1113.
- Felber, A.C., Orlandelli, R.C., Rhoden, S.A., Garcia, A., Costa, A.T., Azevedo, J.L., Pamphile, J.A., 2016. Bioprospecting foliar endophytic fungi of *Vitis labrusca* Linnaeus, Bordô and Concord cv. *Ann. Microbiol.* 66, 765–775.
- Fleet, G.H., 1993. In: CRC Press (Ed.), *Wine Microbiology and Biotechnology*. Harwood Academic Publishers, Chur, Switzerland.
- Fleet, G.H., 2003. Yeast interactions and wine flavour. *Int. J. Food Microbiol.* 86, 11–22.
- Fontaine, M.C., Austerlitz, F., Giraud, T., Labbé, F., Papura, D., Richard-Cervera, S., Delmotte, F., 2013. Genetic signature of a range expansion and leap-frog event after the recent invasion of Europe by the grapevine downy mildew pathogen *Plasmopara viticola*. *Mol. Ecol.* 22, 2771–2786.
- Fontaine, M.C., Labbé, F., Dussert, Y., Delière, L., Richart-Cervera, S., Giraud, T., Delmotte, F., 2021. Europe as a bridgehead in the worldwide invasion history of grapevine downy mildew, *Plasmopara viticola*. *Curr. Biol.* 31, 2155–2166.e4.
- Fort, T., Robin, C., Capdevielle, X., Delière, L., Vacher, C., 2016. Foliar fungal communities strongly differ between habitat patches in a landscape mosaic. *PeerJ* 4, e2656.
- Fort, T., Pauvert, C., Zanne, A.E., Ovaskainen, O., Caignard, T., Barret, M., Compant, S., Hampe, A., Delzon, S., Vacher, C., 2021. Maternal effects shape the seed mycobiome in *Quercus petraea*. *New Phytol.* 230, 1594–1608.
- Friedman, J., Alm, E.J., 2012. Inferring correlation networks from genomic survey data. *PLoS Comput. Biol.* 8, e1002687.
- Friedman, J., Higgins, L.M., Gore, J., 2017. Community structure follows simple assembly rules in microbial microcosms. *Nat. Ecol. Evol.* 1, 1–7.
- Frioux, C., Fremy, E., Trottier, C., Siegel, A., 2018. Scalable and exhaustive screening of metabolic functions carried out by microbial consortia. *Bioinformatics* 34, i934–i943.
- Frioux, C., Singh, D., Korcsmaros, T., Hildebrand, F., 2020. From bag-of-genes to bag-of-genomes: metabolic modelling of communities in the era of metagenome-assembled genomes. *Comput. Struct. Biotechnol. J.* 18, 1722–1734.
- Gadoury, D.M., Cadle-Davidson, L., Wilcox, W.F., Dry, I.B., Seem, R.C., Milgroom, M.G., 2012. Grapevine powdery mildew (*Erysiphe necator*): a fascinating system for the study of the biology, ecology and epidemiology of an obligate biotroph: grapevine powdery mildew. *Mol. Plant Pathol.* 13, 1–16.
- Gambetta, G.A., Herrera, J.C., Dayer, S., Feng, Q., Hochberg, U., Castellarin, S.D., 2020. The physiology of drought stress in grapevine: towards an integrative definition of drought tolerance. *J. Exp. Bot.* 71, 4658–4676.
- Gan, H.M., Szegedi, E., Fersi, R., Chebil, S., Kovács, L., Kawaguchi, A., Hudson, A.O., Burr, T.J., Savka, M.A., 2019. Insight into the microbial co-occurrence and diversity of 73 grapevine (*Vitis vinifera*) crown galls collected across the northern hemisphere. *Front. Microbiol.* 10, 1896.
- Gessler, C., Pertot, I., Perazzolli, M., 2011. *Plasmopara viticola*: a review of knowledge on downy mildew of grapevine and effective disease management. *Phytopathol. Mediterr.* 50, 3–44.
- Ghule, S.B., Sawant, I.S., Sawant, S.D., Saha, S., Devarumath, R.M., 2019. Isolation and identification of three new mycoparasites of *Erysiphe necator* for biological control of grapevine powdery mildew. *Australas. Plant Pathol.* 48, 351–367.
- Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., Egozcue, J.J., 2017. Microbiome datasets are compositional: and this is not optional. *Front. Microbiol.* 8, 2224.



- Gobbi, A., Kyrkou, I., Filippi, E., Ellegaard-Jensen, L., Hansen, L.H., 2020. Seasonal epiphytic microbial dynamics on grapevine leaves under biocontrol and copper fungicide treatments. *Sci. Rep.* 10, 681.
- Gobbi, A., Acedo, A., Imam, N., Santini, R.G., Ortiz-Álvarez, R., Ellegaard-Jensen, L., Belda, I., Hansen, L.H., 2022. A global microbiome survey of vineyard soils highlights the microbial dimension of viticultural terroirs. *Commun. Biol.* 5, 241.
- Gobbin, D., Rumbou, A., Linde, C.C., Gessler, C., 2006. Population genetic structure of *Plasmopara viticola* after 125 years of colonization in European vineyards: Population genetics of *P. viticola* in Europe. *Mol. Plant Pathol.* 7, 519–531.
- González-García, S., Álvarez-Pérez, J.M., Sáenz de Miera, L.E., Cobos, R., Ibañez, A., Díez-Galán, A., Garzón-Jimeno, E., Coque, J.J.R., 2019. Developing tools for evaluating inoculation methods of biocontrol *Streptomyces* sp. strains into grapevine plants. *PLoS One* 14, e0211225.
- Gonze, D., Coyte, K.Z., Lahti, L., Faust, K., 2018. Microbial communities as dynamical systems. *Curr. Opin. Microbiol.* 44, 41–49.
- Gopal, M., Gupta, A., 2016. Microbiome selection could spur next-generation plant breeding strategies. *Front. Microbiol.* 7, 1971.
- Gramaje, D., Armengol, J., 2011. Fungal trunk pathogens in the grapevine propagation process: potential inoculum sources, detection, identification, and management strategies. *Plant Dis.* 95, 1040–1055.
- Gramaje, D., Úrbez-Torres, J.R., Sosnowski, M.R., 2018. Managing grapevine trunk diseases with respect to etiology and epidemiology: current strategies and future prospects. *Plant Dis.* 102, 12–39.
- Gramaje, D., Eichmeier, A., Spetik, M., Carbone, M.J., Bujanda, R., Vallance, J., Rey, P., 2022. Exploring the temporal dynamics of the fungal microbiome in rootstocks, the lesser-known half of the grapevine crop. *J. Fungi* 8, 421.
- Grassi, F., De Lorenzis, G., 2021. Back to the origins: background and perspectives of grapevine domestication. *Int. J. Mol. Sci.* 22, 4518.
- Griggs, R.G., Steenwerth, K.L., Mills, D.A., Cantu, D., Bokulich, N.A., 2021. Sources and assembly of microbial communities in vineyards as a functional component of winegrowing. *Front. Microbiol.* 12, 673810.
- Grube, M., Schmid, F., Berg, G., 2011. Black fungi and associated bacterial communities in the phyllosphere of grapevine. *Fungal Biol.* 115, 978–986.
- Gu, C., Kim, G.B., Kim, W.J., Kim, H.U., Lee, S.Y., 2019. Current status and applications of genome-scale metabolic models. *Genome Biol.* 20, 121.
- Guarnaccia, V., Groenewald, J.Z., Woodhall, J., Armengol, J., Cinelli, T., Eichmeier, A., Ezra, D., Fontaine, F., Gramaje, D., Gutierrez-Aguirregabiria, A., Kaliterna, J., Kiss, L., Laignon, P., Luque, J., Mugnai, L., Naor, V., Raposo, R., Sándor, E., Váczy, K.Z., Crous, P.W., 2018. *Diaporthe* diversity and pathogenicity revealed from a broad survey of grapevine diseases in Europe. *Persoonia - Mol. Phylogeny Evol. Fungi* 40, 135–153.
- Gur, L., Reuveni, M., Cohen, Y., Cadle-Davidson, L., Kisselstein, B., Ovadia, S., Frenkel, O., 2021. Population structure of *Erysiphe necator* on domesticated and wild vines in the Middle East raises questions on the origin of the grapevine powdery mildew pathogen. *Environ. Microbiol.* 23, 6019–6037.
- Gutierrez, A., Grillo, M.A., 2022. Effects of domestication on plant-microbiome interactions. *Plant Cell Physiol.*, 1–13. online.
- Habbadi, K., Duplay, Q., Chapulliot, D., Kerzaon, I., Benkirane, R., Benbouazza, A., Wisniewski-Dyé, F., Lavire, C., Achbani, E.H., Vial, L., 2020. Characterization and phylogenetic diversity of *Allorhizobium vitis* isolated from grapevine in Morocco. *J. Appl. Microbiol.* 128, 828–839.

- Hacquard, S., Spaepen, S., Garrido-Oter, R., Schulze-Lefert, P., 2017. Interplay between innate immunity and the plant microbiota. *Annu. Rev. Phytopathol.* 55, 565–589.
- Haidar, R., Yacoub, A., Vallance, J., Compant, S., Antonielli, L., Saad, A., Habenstein, B., Kauffmann, B., Grélard, A., Loquet, A., Attard, E., Guyoneaud, R., Rey, P., 2021. Bacteria associated with wood tissues of Esca-diseased grapevines: functional diversity and synergy with *Fomitiporia mediterranea* to degrade wood components. *Environ. Microbiol.* 23, 6104–6121.
- Haidar R., Compant S., Antonielli L., Pandin C., Yacoub A., Chrysovergi M., Attard E., Rey P. and Guyoneaud R. *Paenibacillus xylinteritus*, a novel bacterial species isolated from the wood of the grapevine cultivar, Sauvignon Blanc. *Int. J. Syst. Evol. Microbiol.* (submitted).
- Hao, Z., van Tuinen, D., Fayolle, L., Chatagnier, O., Li, X., Chen, B., Gianinazzi, S., Gianinazzi-Pearson, V., 2018. Arbuscular mycorrhiza affects grapevine fanleaf virus transmission by the nematode vector *Xiphinema index*. *Appl. Soil Ecol.* 129, 107–111.
- Heid, C.A., Stevens, J., Livak, K.J., Williams, P.M., 1996. Real time quantitative PCR. *Genome Res.* 6, 986–994.
- Héloir, M.-C., Adrian, M., Brulé, D., Claverie, J., Cordelier, S., Daire, X., Dorey, S., Gauthier, A., Lemaître-Guillier, C., Negrel, J., Trdá, L., Trouvelot, S., Vandelle, E., Poinssot, B., 2019. Recognition of elicitors in grapevine: from MAMP and DAMP perception to induced resistance. *Front. Plant Sci.* 10, 1117.
- Hernandez, M.M., Menéndez, C.M., 2019. Influence of seasonality and management practices on diversity and composition of fungal communities in vineyard soils. *Appl. Soil Ecol.* 135, 113–119.
- Hildebrand, F., Tadeo, R., Voigt, A.Y., Bork, P., Raes, J., 2014. LotuS: an efficient and user-friendly OTU processing pipeline. *Microbiome* 2, 30.
- Hocking, A.D., Leong, S.L., Kazi, B.A., Emmett, R.W., Scott, E.S., 2007. Fungi and mycotoxins in vineyards and grape products. *Int. J. Food Microbiol.* 119, 84–88.
- Hofstetter, V., Buyck, B., Croll, D., Viret, O., Couloux, A., Gindro, K., 2012. What if esca disease of grapevine were not a fungal disease? *Fungal Divers.* 54, 51–67.
- Jaarsveld, W.J., Halleen, F., Bester, M.C., Pierron, R.J., Stempien, E., Mostert, L., 2021. Investigation of *Trichoderma* species colonization of nursery grapevines for improved management of black foot disease. *Pest Manag. Sci.* 77, 397–405.
- Jacoby, R., Peukert, M., Succurro, A., Koprivova, A., Kopriva, S., 2017. The role of soil microorganisms in plant mineral nutrition—current knowledge and future directions. *Front. Plant Sci.* 8, 1617.
- Jacquet, F., Jeuffroy, M.-H., Jouan, J., Le Cadre, E., Litrico, I., Malausa, T., Reboud, X., Huyghe, C., 2022. Pesticide-free agriculture as a new paradigm for research. *Agron. Sustain. Dev.* 42, 8.
- Jakuschkin, B., Fievet, V., Schwaller, L., Fort, T., Robin, C., Vacher, C., 2016. Deciphering the pathobiome: intra- and interkingdom interactions involving the pathogen *Erysiphe alphitoides*. *Microb. Ecol.* 72, 870–880.
- Jara, C., Laurie, V.F., Mas, A., Romero, J., 2016. Microbial terroir in Chilean valleys: diversity of non-conventional yeast. *Front. Microbiol.* 7, 663.
- Jassey, V.E.J., Reczuga, M.K., Zielińska, M., Słowińska, S., Robroek, B.J.M., Mariotte, P., Seppey, C.V.W., Lara, E., Barabach, J., Słowiński, M., Bragazza, L., Chojnicki, B.H., Lamentowicz, M., Mitchell, E.A.D., Buttler, A., 2018. Tipping point in plant–fungal interactions under severe drought causes abrupt rise in peatland ecosystem respiration. *Glob. Chang. Biol.* 24, 972–986.
- Ji, W., Han, K., Cai, Y., Mu, Y., Zhao, L., Zhang, M., Hou, C., Gao, M., Zhao, Q., 2019. Characterization of rhizosphere bacterial community and berry quality of Hutai No.8 (*Vitis vinifera* L.) with different ages, and their relations. *J. Sci. Food Agric.* 99, 4532–4539.

- Joseph, T.A., Shenhav, L., Xavier, J.B., Halperin, E., Peer, I., 2020. Compositional Lotka-Volterra describes microbial dynamics in the simplex. *PLoS Comput. Biol.* 16, e1007917.
- Jürges, G., Kassemeyer, H.-H., Dürrenberger, M., Düggelein, M., Nick, P., 2009. The mode of interaction between *Vitis* and *Plasmopara viticola* Berk. & Curt. Ex de Bary depends on the host species. *Plant Biol.* 11, 886–898.
- Kaminski, T.S., Scheler, O., Garstecki, P., 2016. Droplet microfluidics for microbiology: techniques, applications and challenges. *Lab Chip* 16, 2168–2187.
- Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., Wang, Z., 2019. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* 7, e7359.
- Karp, P.D., Weaver, D., Latendresse, M., 2018. How accurate is automated gap filling of metabolic models? *BMC Syst. Biol.* 12, 73.
- Karp, P.D., Midford, P.E., Billington, R., Kothari, A., Krummenacker, M., Latendresse, M., Ong, W.K., Subhraveti, P., Caspi, R., Fulcher, C., Keseler, I.M., Paley, S.M., 2019. Pathway Tools version 23.0 update: software for pathway/genome informatics and systems biology. *Brief. Bioinform.* 22, 109–126.
- Kerdraon, L., Barret, M., Laval, V., Suffert, F., 2019. Differential dynamics of microbial community networks help identify microorganisms interacting with residue-borne pathogens: the case of *Zymoseptoria tritici* in wheat. *Microbiome* 7, 125.
- Kernaghan, G., Mayerhofer, M., Griffin, A., 2017. Fungal endophytes of wild and hybrid *Vitis* leaves and their potential for vineyard biocontrol. *Can. J. Microbiol.* 63, 583–595.
- Khalifa, M.E., MacDiarmid, R.M., 2021. A mechanically transmitted DNA mycovirus is targeted by the defence machinery of its host, *Botrytis cinerea*. *Viruses* 13, 1315.
- Khodayari, A., Zomorodi, A.R., Liao, J.C., Maranas, C.D., 2014. A kinetic model of *Escherichia coli* core metabolism satisfying multiple sets of mutant flux data. *Metab. Eng.* 25, 50–62.
- Kiss, L., 2003. A review of fungal antagonists of powdery mildews and their potential as biocontrol agents. *Pest Manag. Sci.* 59, 475–483.
- Kleyer, H., Tecon, R., Or, D., 2017. Resolving species level changes in a representative soil bacterial community using microfluidic quantitative PCR. *Front. Microbiol.* 8, 2017.
- Knapp, D.G., Lázár, A., Molnár, A., Vajna, B., Karácsony, Z., Váczy, K.Z., Kovács, G.M., 2021. Above-ground parts of white grapevine *Vitis vinifera* cv. Furmint share core members of the fungal microbiome. *Environ. Microbiol. Rep.* 13, 509–520.
- Kortekamp, A., 1997. *Epicoccum nigrum* Link: a biological control agent of *Plasmopara viticola* (Berk et Curt.) Berl. et De Toni? *Vitis* 36, 215–216.
- Kraus, C., Voegelé, R.T., Fischer, M., 2019. Temporal development of the culturable, endophytic fungal community in healthy grapevine branches and occurrence of GTD-associated fungi. *Microb. Ecol.* 77, 866–876.
- Kuntal, B.K., Gadgil, C., Mande, S.S., 2019. Web-gLV: a web based platform for lotka-volterra based modeling and simulation of microbial populations. *Front. Microbiol.* 10, 288.
- Kurtz, Z.D., Müller, C.L., Miraldi, E.R., Littman, D.R., Blaser, M.J., Bonneau, R.A., 2015. Sparse and compositionally robust inference of microbial ecological networks. *PLoS Comput. Biol.* 11, e1004226.
- La Guerche, S., Chamont, S., Blancard, D., Dubourdieu, D., Darriet, P., 2005. Origin of (–)-geosmin on grapes: on the complementary action of two fungi, *Botrytis cinerea* and *Penicillium expansum*. *Antonie Van Leeuwenhoek* 88, 131–139.
- La Guerche, S., De Senneville, L., Blancard, D., Darriet, P., 2007. Impact of the *Botrytis cinerea* strain and metabolism on (–)-geosmin production by *Penicillium expansum* in grape juice. *Antonie Van Leeuwenhoek* 92, 331–341.

- Labarthe, S., Polizzi, B., Phan, T., Goudon, T., Ribot, M., Laroche, B., 2019. A mathematical model to investigate the key drivers of the biogeography of the colon microbiota. *J. Theor. Biol.* 462, 552–581.
- Lacombe, T., Boursiquot, J.-M., Laucou, V., Di Vecchi-Staraz, M., Péros, J.-P., This, P., 2013. Large-scale parentage analysis in an extended set of grapevine cultivars (*Vitis vinifera* L.). *Theor. Appl. Genet.* 126, 401–414.
- Lade, S.B., Štraus, D., Oliva, J., 2022. Variation in fungal community in grapevine (*Vitis vinifera*) nursery stock depends on nursery, variety and rootstock. *J. Fungi* 8, 47.
- Lagier, J.-C., Hugon, P., Khelaïfia, S., Fournier, P.-E., La Scola, B., Raoult, D., 2015. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin. Microbiol. Rev.* 28, 237–264.
- Lagier, J.-C., Khelaïfia, S., Alou, M.T., Ndongo, S., Dione, N., Hugon, P., Caputo, A., Cadoret, F., Traore, S.I., Seck, E.H., Dubourg, G., Durand, G., Mourembou, G., Guilhot, E., Togo, A., Bellali, S., Bachar, D., Cassir, N., Bittar, F., Delerce, J., Mailhe, M., Ricaboni, D., Bilén, M., Dangui Niekou, N.P.M., Dia Badiane, N.M., Valles, C., Mouelhi, D., Diop, K., Million, M., Musso, D., Abrahão, J., Azhar, E.I., Bibi, F., Yasir, M., Diallo, A., Sokhna, C., Djossou, F., Vitton, V., Robert, C., Rolain, J.M., La Scola, B., Fournier, P.-E., Levasseur, A., Raoult, D., 2016. Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat. Microbiol.* 1, 1–8.
- Lagier, J.-C., Dubourg, G., Million, M., Cadoret, F., Bilén, M., Fenollar, F., Levasseur, A., Rolain, J.-M., Fournier, P.-E., Raoult, D., 2018. Culturing the human microbiota and culturomics. *Nat. Rev. Microbiol.* 16, 540–550.
- Laimer, M., Lemaire, O., Herrbach, E., Goldschmidt, V., Minafra, A., Bianco, P., Wetzell, T., 2009. Resistance to viruses, phytoplasmas and their vectors in the grapevine in Europe: a review. *J. Plant Pathol.* 91, 7–23.
- Legras, J.-L., Merdinoglu, D., Cornuet, J.-M., Karst, F., 2007. Bread, beer and wine: *Saccharomyces cerevisiae* diversity reflects human history. *Mol. Ecol.* 16, 2091–2102.
- Lemos, L.N., Mendes, L.W., Baldrian, P., Pylro, V.S., 2021. Genome-resolved metagenomics is essential for unlocking the microbial black box of the soil. *Trends Microbiol.* 29, 279–282.
- Levy, A., Salas Gonzalez, I., Mittelviefhaus, M., Clingenpeel, S., Herrera Paredes, S., Miao, J., Wang, K., Devescovi, G., Stillman, K., Monteiro, F., Rangel Alvarez, B., Lundberg, D.S., Lu, T.-Y., Lebeis, S., Jin, Z., McDonald, M., Klein, A.P., Feltscher, M.E., Rio, T.G., Grant, S.R., Doty, S.L., Ley, R.E., Zhao, B., Venturi, V., Pelletier, D.A., Vorholt, J.A., Tringe, S.G., Woyke, T., Dangl, J.L., 2018. Genomic features of bacterial adaptation to plants. *Nat. Genet.* 50, 138–150.
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31, 1674–1676.
- Li, R., Yang, S., Lin, M., Guo, S., Han, X., Ren, M., Du, L., Song, Y., You, Y., Zhan, J., Huang, W., 2022. The biogeography of fungal communities across different chinese wine-producing regions associated with environmental factors and spontaneous fermentation performance. *Front. Microbiol.* 12, 636639.
- Likar, M., Stres, B., Rusjan, D., Potisek, M., Regvar, M., 2017. Ecological and conventional viticulture gives rise to distinct fungal and bacterial microbial communities in vineyard soils. *Appl. Soil Ecol.* 113, 86–95.
- Liu, D., Howell, K., 2021. Community succession of the grapevine fungal microbiome in the annual growth cycle. *Environ. Microbiol.* 23, 1842–1857.
- Liu, D., Zhang, P., Chen, D., Howell, K., 2019. From the vineyard to the winery: how microbial ecology drives regional distinctiveness of wine. *Front. Microbiol.* 10, 2679.

- Liu, Z., Ma, A., Mathé, E., Merling, M., Ma, Q., Liu, B., 2021. Network analyses in microbiome based on high-throughput multi-omics data. *Brief. Bioinform.* 22 (2), 1639–1655.
- Lleixa, J., Kioroglou, D., Mas, A., Portillo, M., 2018. Microbiome dynamics during spontaneous fermentations of sound grapes in comparison with sour rot and *Botrytis* infected grapes. *Int. J. Food Microbiol.* 281, 36–46.
- Lo, C., Marculescu, R., 2017. MPLasso: Inferring microbial association networks using prior microbial knowledge. *PLoS Comput. Biol.* 13, e1005915.
- Loqman, S., Barka, E.A., Clément, C., Ouhdouch, Y., 2009. Antagonistic actinomycetes from Moroccan soil to control the grapevine gray mold. *World J. Microbiol. Biotechnol.* 25, 81–91.
- Lou, J., Yang, L., Wang, H., Wu, L., Xu, J., 2018. Assessing soil bacterial community and dynamics by integrated high-throughput absolute abundance quantification. *PeerJ* 6, e4514.
- Louca, S., Parfrey, L.W., Doebeli, M., 2016. Decoupling function and taxonomy in the global ocean microbiome. *Science* 353, 1272–1277.
- Lyu, D., Msimbira, L.A., Nazari, M., Antar, M., Pagé, A., Shah, A., Monjezi, N., Zajonc, J., Tanney, C.A.S., Backer, R., Smith, D.L., 2021. The coevolution of plants and microbes underpins sustainable agriculture. *Microorganisms* 9, 1036.
- Magnin-Robert, M., Trotel-Aziz, P., Quantinet, D., Biagianni, S., Aziz, A., 2007. Biological control of *Botrytis cinerea* by selected grapevine-associated bacteria and stimulation of chitinase and  $\beta$ -1,3 glucanase activities under field conditions. *Eur. J. Plant Pathol.* 118, 43–57.
- Mahadevan, R., Edwards, J.S., Doyle, F.J., 2002. Dynamic flux balance analysis of diauxic growth in *Escherichia coli*. *Biophys. J.* 83, 1331–1340.
- Maher, N., Piot, J., Bastien, S., Vallance, J., Rey, P., Guérin-Dubrana, L., 2012. Wood necrosis in ESCA-affected vines: types, relationships and possible links with foliar symptom expression. *J. Int. Sci. Vigne Vin* 46, 15–27.
- Majer, P., Hideg, É., 2012. Developmental stage is an important factor that determines the antioxidant responses of young and old grapevine leaves under UV irradiation in a green-house. *Plant Physiol. Biochem.* 50, 15–23.
- Maliogka, V.I., Martelli, G.P., Fuchs, M., Katis, N.I., 2015. Control of viruses infecting grapevine. In: *Advances in Virus Research*. Elsevier, pp. 175–227.
- Marais, A., Faure, C., Comont, G., Candresse, T., Stempien, E., Corio-Costet, M.-F., 2021. Characterization of the mycovirome of the phytopathogenic fungus, *Neofusicoccum parvum*. *Viruses* 13, 375.
- Marasco, R., Rolli, E., Fusi, M., Michoud, G., Daffonchio, D., 2018. Grapevine rootstocks shape underground bacterial microbiome and networking but not potential functionality. *Microbiome* 6, 3.
- Marchesi, J.R., Ravel, J., 2015. The vocabulary of microbiome research: a proposal. *Microbiome* 3, 31.
- Mardanov, A.V., Chizhevskaya, E.P., Lazarev, A.M., Rakitin, A.L., Beletsky, A.V., Chebotar, V.K., Ravin, N.V., 2019. Complete genome sequences of endophytic bacilli isolated from grapevine plants. *Microbiol. Resour. Announc.* 8, e01265–19.
- Marín, O., González, B., Poupin, M.J., 2021. From microbial dynamics to functionality in the rhizosphere: a systematic review of the opportunities with synthetic microbial communities. *Front. Plant Sci.* 12, 650609.
- Marino, S., Baxter, N.T., Huffnagle, G.B., Petrosino, J.F., Schloss, P.D., 2014. Mathematical modeling of primary succession of murine intestinal microbiota. *Proc. Natl. Acad. Sci. U. S. A.* 111, 439–444.
- Marsit, S., Dequin, S., 2015. Diversity and adaptive evolution of *Saccharomyces* wine yeast: a review. *FEMS Yeast Res.* 15, fov067.

- Martelli, G.P., 1999. Proceedings of the Mediterranean Network on Grapevine Closteroviruses 1992-1997 and the Viroses and Virus-Like Diseases of the Grapevine. Centre international de hautes études agronomiques (méditerranéennes).
- Martini, M., Musetti, R., Grisan, S., Polizzotto, R., Borselli, S., Pavan, F., Osler, R., 2009. DNA-dependent detection of the grapevine fungal endophytes *Aureobasidium pullulans* and *Epicoccum nigrum*. *Plant Dis.* 93, 993–998.
- Martins, G., Miot-Sertier, C., Lauga, B., Claisse, O., Lonvaud-Funel, A., Soulas, G., Masneuf-Pomarède, I., 2012. Grape berry bacterial microbiota: impact of the ripening process and the farming system. *Int. J. Food Microbiol.* 158, 93–100.
- Martins, G., Vallance, J., Mercier, A., Albertin, W., Stamatopoulos, P., Rey, P., Lonvaud, A., Masneuf-Pomarède, I., 2014. Influence of the farming system on the epiphytic yeasts and yeast-like fungi colonizing grape berries during the ripening process. *Int. J. Food Microbiol.* 177, 21–28.
- Martins, G.E., Casini, C., Da Costa, J.-P., Geny, L., Lonvaud, A., Masneuf-Pomarède, I., 2020. Correlation between water activity (aw) and microbial epiphytic communities associated with grapes berries. *OENO One* 54, 49–61.
- Mendoza, S.N., Cañón, P.M., Contreras, Á., Ribbeck, M., Agosín, E., 2017. Genome-scale reconstruction of the metabolic network in *Oenococcus oeni* to assess wine malolactic fermentation. *Front. Microbiol.* 8, 534.
- Merdinoglu, D., Schneider, C., Prado, E., Wiedemann-Merdinoglu, S., Mestre, P., 2018. Breeding for durable resistance to downy and powdery mildew in grapevine. *OENO One* 52, 203–209.
- Miotto-Vilanova, L., Jacquard, C., Courteaux, B., Wortham, L., Michel, J., Clément, C., Barka, E.A., Sanchez, L., 2016. *Burkholderia phytofirmans* PsJN confers grapevine resistance against *Botrytis cinerea* via a direct antimicrobial effect combined with a better resource mobilization. *Front. Plant Sci.* 7, 1236.
- Miura, T., Sánchez, R., Castañeda, L.E., Godoy, K., Barbosa, O., 2017. Is microbial terroir related to geographic distance between vineyards? Microbial terroir related to geographic distance. *Environ. Microbiol. Rep.* 9, 742–749.
- Miura, T., Sánchez, R., Castañeda, L.E., Godoy, K., Barbosa, O., 2019. Shared and unique features of bacterial communities in native forest and vineyard phyllosphere. *Ecol. Evol.* 9, 3295–3305.
- Molitor, D., Beyer, M., 2014. Epidemiology, identification and disease management of grape black rot and potentially useful metabolites of black rot pathogens for industrial applications—a review. *Ann. Appl. Biol.* 165, 305–317.
- Momeni, B., Xie, L., Shou, W., 2017. Lotka-Volterra pairwise modeling fails to capture diverse pairwise microbial interactions. *Elife* 6, e25051.
- Mondello, V., Songy, A., Battiston, E., Pinto, C., Coppin, C., Trotel-Aziz, P., Clément, C., Mugnai, L., Fontaine, F., 2018. Grapevine trunk diseases: a review of fifteen years of trials for their control with chemicals and biocontrol agents. *Plant Dis.* 102, 1189–1217.
- Morales-Cedeño, L.R., Ma del Orozco-Mosqueda, C., Loeza-Lara, P.D., Parra-Cota, F.I., de los Santos-Villalobos, S., Santoyo, G., 2021. Plant growth-promoting bacterial endophytes as biocontrol agents of pre- and post-harvest diseases: fundamentals, methods of application and future perspectives. *Microbiol. Res.* 242, 126612.
- Morgan, H.H., du Toit, M., Setati, M.E., 2017. The grapevine and wine microbiome: insights from high-throughput amplicon sequencing. *Front. Microbiol.* 8, 820.
- Mortimer, R., Polsinelli, M., 1999. On the origins of wine yeast. *Res. Microbiol.* 150 (3), 199–204.
- Mourad, E.F., Sarhan, M.S., Daanaa, H.-S.A., Abdou, M., Morsi, A.T., Abdelfadeel, M.R., Elsayey, H., Nemr, R., El-Tahan, M., Hamza, M.A., Abbas, M., Youssef, H.H., Abdelhadi, A.A., Amer, W.M., Fayez, M., Ruppel, S., Hegazi, N.A., 2018. Plant

- materials are sustainable substrates supporting new technologies of plant-only-based culture media for *in vitro* Culturing of the plant microbiota. *Microbes Environ.* 33 (1), 40–49. adypub, ME17135.
- Mugnai, L., Graniti, A., Surico, G., 1999. Esca (black measles) and brown wood-streaking: two old and elusive diseases of grapevines. *Plant Dis.* 83, 404–418.
- Musetti, R., Vecchione, A., Stringher, L., Borselli, S., Zulini, L., Marzani, C., D'Ambrosio, M., di Toppi, L.S., Pertot, I., 2006. Inhibition of sporulation and ultrastructural alterations of grapevine downy mildew by the endophytic fungus *alternaria alternata*. *Phytopathology* 96, 689–698.
- Myles, S., Boyko, A.R., Owens, C.L., Brown, P.J., Grassi, F., Aradhya, M.K., Prins, B., Reynolds, A., Chia, J.-M., Ware, D., Bustamante, C.D., Buckler, E.S., 2011. Genetic structure and domestication history of the grape. *Proc. Natl. Acad. Sci. U. S. A.* 108, 3530–3535.
- Nai, C., Meyer, V., 2018. From axenic to mixed cultures: technological advances accelerating a paradigm shift in microbiology. *Trends Microbiol.* 26, 538–554.
- Narduzzi-Wicht, B., Jermini, M., Gessler, C., Broggin, G.A.L., 2014. Microsatellite markers for population studies of the ascomycete *Phyllosticta ampellicida*, the pathogen causing grape black rot. *Phytopathol. Mediterr.* 53, 470–479.
- Nayfach, S., Roux, S., Seshadri, R., Udway, D., Varghese, N., Schulz, F., Wu, D., Paez-Espino, D., Chen, I.-M., Huntemann, M., Palaniappan, K., Ladau, J., Mukherjee, S., Reddy, T.B.K., Nielsen, T., Kirton, E., Faria, J.P., Edirisinghe, J.N., Henry, C.S., Jungbluth, S.P., Chivian, D., Dehal, P., Wood-Charlson, E.M., Arkin, A.P., Tringe, S.G., Visel, A., Abreu, H., Acinas, S.G., Allen, E., Allen, M.A., Andersen, G., Anesio, A.M., Attwood, G., Avila-Magaña, V., Badis, Y., Bailey, J., Baker, B., Baldrian, P., Barton, H.A., Beck, D.A.C., Becraft, E.D., Beller, H.R., Beman, J.M., Bernier-Latmani, R., Berry, T.D., Bertagnolli, A., Bertilsson, S., Bhatnagar, J.M., Bird, J.T., Blumer-Schuette, S.E., Bohannan, B., Borton, M.A., Brady, A., Brawley, S.H., Brodie, J., Brown, S., Brum, J.R., Brune, A., Bryant, D.A., Buchan, A., Buckley, D.H., Buongiorno, J., Cadillo-Quiroz, H., Caffrey, S.M., Campbell, A.N., Campbell, B., Carr, S., Carroll, J., Cary, S.C., Cates, A.M., Cattolico, R.A., Cavicchioli, R., Chistoserdova, L., Coleman, M.L., Constant, P., Conway, J.M., Cormack, W.P.M., Crowe, S., Crump, B., Currie, C., Daly, R., Denev, V., Denman, S.E., Desta, A., Dionisi, H., Dodsworth, J., Dombrowski, N., Donohue, T., Dopson, M., Driscoll, T., Dunfield, P., Dupont, C.L., Dynarski, K.A., Edgcomb, V., Edwards, E.A., Elshahed, M.S., Figueroa, I., Flood, B., Fortney, N., Fortunato, C.S., Francis, C., Gachon, C.M.M., Garcia, S.L., Gazitua, M.C., Gentry, T., Gerwick, L., Gharechahi, J., Girguis, P., Gladden, J., Gradoville, M., Grasby, S.E., Gravier, K., Grettenberger, C.L., Gruninger, R.J., Guo, J., Habteselassie, M.Y., Hallam, S.J., Hatzepichler, R., Hausmann, B., Hazen, T.C., Hedlund, B., Henny, C., Herfort, L., Hernandez, M., Hershey, O.S., Hess, M., Hollister, E.B., Hug, L.A., Hunt, D., Jansson, J., Jarett, J., Kadnikov, V.V., Kelly, C., Kelly, R., Kelly, W., Kerfeld, C.A., Kimbrel, J., Klassen, J.L., Konstantinidis, K.T., Lee, L.L., Li, W.-J., Loder, A.J., Loy, A., Lozada, M., MacGregor, B., Magnabosco, C., da Silva, A.M., McKay, R.M., McMahon, K., McSweeney, C.S., Medina, M., Meredith, L., Mizzi, J., Mock, T., Momper, L., Moran, M.A., Morgan-Lang, C., Moser, D., Muiyzer, G., Myrold, D., Nash, M., Nesbø, C.L., Neumann, A.P., Neumann, R.B., Noguera, D., Northen, T., Norton, J., Nowinski, B., Nüsslein, K., O'Malley, M.A., Oliveira, R.S., de Oliveira, V.M., Onstott, T., Osvatic, J., Ouyang, Y., Pachiadaki, M., Parnell, J., Partida-Martinez, L.P., Peay, K.G., Pelletier, D., Peng, X., Pester, M., Pett-Ridge, J., Peura, S., Pjevac, P., Plominsky, A.M., Poehlein, A., Pope, P.B., Ravin, N., Redmond, M.C., Reiss, R., Rich, V.,



- Rinke, C., Rodrigues, J.L.M., Rossmassler, K., Sackett, J., Salekdeh, G.H., Saleska, S., Scarborough, M., Schachtman, D., Schadt, C.W., Schrenk, M., Szczyrba, A., Sengupta, A., Setubal, J.C., Shade, A., Sharp, C., Sherman, D.H., Shubenkova, O.V., Sierra-Garcia, I.N., Simister, R., Simon, H., Sjöling, S., Slonczewski, J., de Souza, R.S.C., Spear, J.R., Stegen, J.C., Stepanauskas, R., Stewart, F., Suen, G., Sullivan, M., Sumner, D., Swan, B.K., Swingle, W., Tarn, J., Taylor, G.T., Teeling, H., Tekere, M., Teske, A., Thomas, T., Thrash, C., Tiedje, J., Ting, C.S., Tully, B., Tyson, G., Ulloa, O., Valentine, D.L., Goethem, M.W.V., VanderGheynst, J., Verbeke, T.J., Vollmers, J., Vuillemin, A., Waldo, N.B., Walsh, D.A., Weimer, B.C., Whitman, T., van der Wielen, P., Wilkins, M., Williams, T.J., Woodcroft, B., Woollet, J., Wrighton, K., Ye, J., Young, E.B., Youssef, N.H., Yu, F.B., Zemska, T.I., Ziels, R., Woyke, T., Mouncey, N.J., Ivanova, N.N., Kyrpides, N.C., Eloe-Fadrosh, E.A., 2020. A genomic catalog of Earth's microbiomes. *Nat. Biotechnol.* 39, 499–509.
- Nearing, J.T., Douglas, G.M., Hayes, M.G., MacDonald, J., Desai, D.K., Allward, N., Jones, C.M.A., Wright, R.J., Dhanani, A.S., Comeau, A.M., Langille, M.G.I., 2022. Microbiome differential abundance methods produce different results across 38 datasets. *Nat. Commun.* 13, 342.
- Nemergut, D.R., Schmidt, S.K., Fukami, T., O'Neill, S.P., Bilinski, T.M., Stanish, L.F., Knelman, J.E., Darcy, J.L., Lynch, R.C., Wickey, P., Ferrenberg, S., 2013. Patterns and processes of microbial community assembly. *Microbiol. Mol. Biol. Rev.* 77, 342–356.
- Nerva, L., Turina, M., Zanzotto, A., Gardiman, M., Gaiotti, F., Gambino, G., Chitarra, W., 2019. Isolation, molecular characterization and virome analysis of culturable wood fungal endophytes in esca symptomatic and asymptomatic grapevine plants. *Environ. Microbiol.* 21, 2886–2904.
- Nerva, L., Garcia, J.F., Favaretto, F., Giudice, G., Moffa, L., Sandrini, M., Cantu, D., Zanzotto, A., Gardiman, M., Velasco, R., Gambino, G., Chitarra, W., 2022. The hidden world within plants: metatranscriptomics unveils the complexity of wood microbiomes. *J. Exp. Bot.* 73, 2682–2697.
- Nguyen, N.H., Song, Z., Bates, S.T., Branco, S., Tedersoo, L., Menke, J., Schilling, J.S., Kennedy, P.G., 2016. FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol.* 20, 241–248.
- Nidelet, T., Brial, P., Camarasa, C., Dequin, S., 2016. Diversity of flux distribution in central carbon metabolism of *S. cerevisiae* strains from diverse environments. *Microb. Cell Fact.* 15, 58.
- Ning, D., Deng, Y., Tiedje, J.M., Zhou, J., 2019. A general framework for quantitatively assessing ecological stochasticity. *Proc. Natl. Acad. Sci. U. S. A.* 116, 16892–16898.
- Nisiotou, A.A., Nychas, G.J., 2007. Yeast populations residing on healthy or botrytis-infected grapes from a vineyard in Attica, Greece. *Appl. Environ. Microbiol.* 73 (8), 2765–2768.
- Noman, M., Ahmed, U., Ijaz, U., Shahid, M., Azizullah, Li, D., Manzoor, I., Song, F., 2021. Plant–microbiome crosstalk: dawning from composition and assembly of microbial community to improvement of disease resilience in plants. *Int. J. Mol. Sci.* 22, 6852.
- Novello, G., Gamalero, E., Bona, E., Boatti, L., Mignone, F., Massa, N., Cesaro, P., Lingua, G., Berta, G., 2017. The rhizosphere bacterial microbiota of *Vitis vinifera* cv. Pinot noir in an integrated pest management vineyard. *Front. Microbiol.* 8, 1528.
- Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P.A., 2017. metaSPAdes: a new versatile metagenomic assembler. *Genome Res.* 27, 824–834.
- Ocete, R., Armendariz, I., Cantos, M., Álvarez, D., Azcón, R., 2015. Ecological characterization of wild grapevine habitats focused on arbuscular mycorrhizal symbiosis. *Vitis* 54, 207–211.

- Ofek-Lazar, M., Gur, Y., Ben-Moshe, S., Sharon, O., Kosman, E., Mochli, E., Sharon, A., 2016. Diversity of fungal endophytes in recent and ancient wheat ancestors *Triticum dicoccoides* and *Aegilops sharonensis*. *FEMS Microbiol. Ecol.* 92, fiw152.
- Oliveira, M., Arenas, M., Lage, O., Cunha, M., Amorim, M.I., 2018. Epiphytic fungal community in *Vitis vinifera* of the Portuguese wine regions. *Lett. Appl. Microbiol.* 66, 93–102.
- Oliver, J.E., Fuchs, M.F., 2011. Fanleaf Degeneration/Decline Disease of Grapevines. Cornell University and the New York State IPM Program, Ithaca, pp. 1–3.
- Oren, A., Garrity, G.M., 2021. Valid publication of the names of forty-two phyla of prokaryotes. *Int. J. Syst. Evol. Microbiol.* 71 (10), 005056.
- Orth, J.D., Thiele, I., Palsson, B.Ø., 2010. What is flux balance analysis? *Nat. Biotechnol.* 28, 245–248.
- Ouadi, L., Bruez, E., Bastien, S., Vallance, J., Lecomte, P., Domec, J.-C., Rey, P., 2019. Ecophysiological impacts of Esca, a devastating grapevine trunk disease, on *Vitis vinifera* L. *PLoS One* 14, e0222586.
- Ouadi, L., Bruez, E., Bastien, S., Yacoub, A., Coppin, C., Guérin-Dubrana, L., Fontaine, F., Domec, J.-C., Rey, P., 2021. Sap flow disruption in grapevine is the early signal predicting the structural, functional, and genetic responses to esca disease. *Front. Plant Sci.* 12, 695846.
- Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., Roslin, T., Abrego, N., 2017. How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecol. Lett.* 20, 561–576.
- Pacifico, D., Squartini, A., Crucitti, D., Barizza, E., Lo Schiavo, F., Muresu, R., Carimi, F., Zottini, M., 2019. The role of the endophytic microbiome in the grapevine response to environmental triggers. *Front. Plant Sci.* 10, 1256.
- Pandey, B., Naidu, R.A., Grove, G.G., 2018. Detection and analysis of mycovirus-related RNA viruses from grape powdery mildew fungus *Erysiphe necator*. *Arch. Virol.* 163, 1019–1030.
- Panstruga, R., Kuhn, H., 2019. Mutual interplay between phytopathogenic powdery mildew fungi and other microorganisms. *Mol. Plant Pathol.* 20, 463–470.
- Paolinelli, M., Escoriaza, G., Cesari, C., Garcia-Lampasona, S., Hernandez-Martinez, R., 2022. Characterization of grapevine wood microbiome through a metatranscriptomic approach. *Microb. Ecol.* 83, 658–668.
- Pauvert, C., Buée, M., Laval, V., Edel-Hermann, V., Fauchery, L., Gautier, A., Lesur, I., Vallance, J., Vacher, C., 2019. Bioinformatics matters: the accuracy of plant and soil fungal community data is highly dependent on the metabarcoding pipeline. *Fungal Ecol.* 41, 23–33.
- Pauvert, C., Fort, T., Calonnec, A., Faivre d'Arcier, J., Chancerel, E., Massot, M., Chiquet, J., Robin, S., Bohan, D.A., Vallance, J., Vacher, C., 2020. Microbial association networks give relevant insights into plant pathobiomes. *bioRxiv*. 2020.02.21.958033.
- Pearson, M.N., Bailey, A.M., 2013. Viruses of *Botrytis*. In: *Advances in Virus Research*. Elsevier, pp. 249–272.
- Perazzolli, M., Antonielli, L., Storari, M., Puopolo, G., Pancher, M., Giovannini, O., Pindo, M., Pertot, I., 2014. Resilience of the natural phyllosphere microbiota of the grapevine to chemical and biological pesticides. *Appl. Environ. Microbiol.* 80, 3585–3596.
- Perera, D., Savocchia, S., Prenzler, P.D., Thomson, P.C., Steel, C.C., 2021. Occurrence of fumonisin-producing black aspergilli in Australian wine grapes: effects of temperature and water activity on fumonisin production by *A. niger* and *A. welwitschiae*. *Mycotoxin Res.* 37, 327–339.

- Peressotti, E., Wiedemann-Merdinoglu, S., Delmotte, F., Bellin, D., Di Gaspero, G., Testolin, R., Merdinoglu, D., Mestre, P., 2010. Breakdown of resistance to grapevine downy mildew upon limited deployment of a resistant variety. *BMC Plant Biol.* 10, 147.
- Pérez-Valera, E., Verdú, M., Navarro-Cano, J.A., Goberna, M., 2020. Soil microbiome drives the recovery of ecosystem functions after fire. *Soil Biol. Biochem.* 149, 107948.
- Pertot, I., Caffi, T., Rossi, V., Mugnai, L., Hoffmann, C., Grando, M.S., Gary, C., Lafond, D., Duso, C., Thiery, D., Mazzoni, V., Anfora, G., 2017. A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for the implementation of IPM in viticulture. *Crop Prot.* 97, 70–84. Pesticide use and risk reduction with IPM.
- Philippe, C., Krupovic, M., Jaomanjaka, F., Claisse, O., Petrel, M., le Marrec, C., 2018. Bacteriophage GC1, a novel tectivirus infecting *Gluconobacter Cerinus*, an acetic acid bacterium associated with wine-making. *Viruses* 10, 39.
- Pilar Martínez-Diz, M., Díaz-Losada, E., Andrés-Sodupe, M., Bujanda, R., Maldonado-González, M.M., Ojeda, S., Yacoub, A., Rey, P., Gramaje, D., 2021. Field evaluation of biocontrol agents against black-foot and petri diseases of grapevine. *Pest Manag. Sci.* 77, 697–708.
- Pinto, C., Pinho, D., Sousa, S., Pinheiro, M., Egas, C., Gomes, A.C., 2014. Unravelling the diversity of grapevine microbiome. *PLoS One* 9, e85622.
- Pirrello, C., Mizzotti, C., Tomazetti, T.C., Colombo, M., Bettinelli, P., Prodorutti, D., Peressotti, E., Zulini, L., Stefanini, M., Angeli, G., Masiero, S., Welter, L.J., Hausmann, L., Vezzulli, S., 2019. Emergent ascomycetes in viticulture: an interdisciplinary overview. *Front. Plant Sci.* 10, 1394.
- Poudel, R., Jumpponen, A., Schlatter, D.C., Paulitz, T.C., Gardener, B.B.M., Kinkel, L.L., Garrett, K.A., 2016. Microbiome networks: a systems framework for identifying candidate microbial assemblages for disease management. *Phytopathology* 106, 1083–1096.
- Prodhomme, D., Valls Fonayet, J., Hévin, C., Franc, C., Hilbert, G., de Revel, G., Richard, T., Ollat, N., Cookson, S.J., 2019. Metabolite profiling during graft union formation reveals the reprogramming of primary metabolism and the induction of stilbene synthesis at the graft interface in grapevine. *BMC Plant Biol.* 19, 599.
- Props, R., Kerckhof, F.-M., Rubbens, P., De Vrieze, J., Hernandez Sanabria, E., Waegeman, W., Monsieus, P., Hammes, F., Boon, N., 2017. Absolute quantification of microbial taxon abundances. *ISME J.* 11, 584–587.
- Prosser, J.I., Bohannan, B.J.M., Curtis, T.P., Ellis, R.J., Firestone, M.K., Freckleton, R.P., Green, J.L., Green, L.E., Killham, K., Lennon, J.J., Osborn, A.M., Solan, M., van der Gast, C.J., Young, J.P.W., 2007. The role of ecological theory in microbial ecology. *Nat. Rev. Microbiol.* 5, 384–392.
- Puig-Pujol, A., Ferrando, N., Capdevila, F., Ocete, R., Revilla, E., 2016. Yeast biodiversity from *Vitis vinifera* L., subsp. *sylvestris* (Gmelin) Hegi to face up the oenological consequences of climate change. *BIO Web Conf.* 7, 02026.
- Qiu, W., Feechan, A., Dry, I., 2015. Current understanding of grapevine defense mechanisms against the biotrophic fungus (*Erysiphe necator*), the causal agent of powdery mildew disease. *Hortic. Res.* 2, 15020.
- Quince, C., Walker, A.W., Simpson, J.T., Loman, N.J., Segata, N., 2017. Shotgun metagenomics, from sampling to analysis. *Nat. Biotechnol.* 35, 833–844.
- Radić, T., Likar, M., Hančević, K., Regvar, M., Čarija, M., Zdunić, G., 2021. Root-associated community composition and co-occurrence patterns of fungi in wild grapevine. *Fungal Ecol.* 50, 101034.

- Raguideau, S., Plancade, S., Pons, N., Leclerc, M., Laroche, B., 2016. Inferring aggregated functional traits from metagenomic data using constrained non-negative matrix factorization: application to fiber degradation in the human gut microbiota. *PLoS Comput. Biol.* 12, e1005252.
- Raspor, P., Miklič-Milek, D., Avbelj, M., Čadež, N., 2010. *Food Technol. Biotechnol.* 48, 336–343.
- Rathnayake, R.M.S.P., Savocchia, S., Schmidtke, L.M., Steel, C.C., 2018. Characterisation of *Aureobasidium pullulans* isolates from *Vitis vinifera* and potential biocontrol activity for the management of bitter rot of grapes. *Eur. J. Plant Pathol.* 151, 593–611.
- Remien, C.H., Eckwright, M.J., Ridenhour, B.J., 2021. Structural identifiability of the generalized Lotka–Volterra model for microbiome studies. *R. Soc. Open Sci.* 8, 201378.
- Rezgui, A., Ben Ghnaya-Chakroun, A., Vallance, J., Bruez, E., Hajlaoui, M.R., Sadfi-Zouaoui, N., Rey, P., 2016. Endophytic bacteria with antagonistic traits inhabit the wood tissues of grapevines from Tunisian vineyards. *Biol. Control* 99, 28–37.
- Rijavec, T., Lapanje, A., 2016. Hydrogen cyanide in the rhizosphere: not suppressing plant pathogens, but rather regulating availability of phosphate. *Front. Microbiol.* 7, 1785.
- Rillig, M.C., Antonovics, J., Caruso, T., Lehmann, A., Powell, J.R., Veresoglou, S.D., Verbruggen, E., 2015. Interchange of entire communities: microbial community coalescence. *Trends Ecol. Evol.* 30, 470–476.
- Rinaldi, P.A., Paffetti, D., Comparini, C., Broggin, G.A.L., Gessler, C., Mugnai, L., 2017. Genetic variability of *Phyllosticta ampellicida*, the agent of black rot disease of grapevine. *Phytopathology* 107, 1406–1416.
- Riva, V., Mapelli, F., Bagnasco, A., Mengoni, A., Borin, S., 2022. A Meta-analysis approach to defining the culturable core of plant endophytic bacterial communities. *Appl. Environ. Microbiol.* 88, e02537-21.
- Rivas, G.A., Semorile, L., Delfederico, L., 2022. Microbial diversity of the soil, rhizosphere and wine from an emerging wine-producing region of Argentina. *LWT* 153, 112429.
- Rizaludin, M.S., Stopnisek, N., Raaijmakers, J.M., Garbeva, P., 2021. The chemistry of stress: understanding the ‘cry for help’ of plant roots. *Metabolites* 11, 357.
- Rondot, Y., Reineke, A., 2019. Endophytic *Beauveria bassiana* activates expression of defence genes in grapevine and prevents infections by grapevine downy mildew *Plasmopara viticola*. *Plant Pathol.* 68, 1719–1731.
- Roossinck, M.J., 2010. Lifestyles of plant viruses. *Philos. Trans. R. Soc. B Biol. Sci.* 365, 1899–1905.
- Ross, T., McMeekin, T.A., 1994. Predictive microbiology. *Int. J. Food Microbiol.* 23, 241–264. Special Issue Predictive Modelling.
- Röttgers, L., Faust, K., 2018. From hairballs to hypotheses—biological insights from microbial networks. *FEMS Microbiol. Rev.* 42, 761–780.
- Rouxel, M., Mestre, P., Comont, G., Lehman, B.L., Schilder, A., Delmotte, F., 2013. Phylogenetic and experimental evidence for host-specialized cryptic species in a biotrophic oomycete. *New Phytol.* 197, 251–263.
- Rouxel, M., Mestre, P., Baudoin, A., Carisse, O., Delière, L., Ellis, M.A., Gadoury, D., Lu, J., Nita, M., Richard-Cervera, S., Schilder, A., Wise, A., Delmotte, F., 2014. Geographic distribution of cryptic species of *Plasmopara viticola* causing downy mildew on wild and cultivated grape in eastern North America. *Phytopathology* 104, 692–701.
- Rowhani, A., Uyemoto, J.K., Golino, D.A., Daubert, S.D., Al Rwahnih, M., 2017. Viruses involved in graft incompatibility and decline. In: Meng, B., Martelli, G.P., Golino, D.A., Fuchs, M. (Eds.). *Grapevine Viruses: Molecular Biology, Diagnostics and Management*. Springer International Publishing, Cham, pp. 289–302.
- Ruiz-Padilla, A., Rodríguez-Romero, J., Gómez-Cid, I., Pacifico, D., Ayllón, M.A., 2021. Novel mycoviruses discovered in the mycovirome of a necrotrophic fungus. *MBio* 12, e03705–e03720.

- Saikkonen, K., Nissinen, R., Helander, M., 2020. Toward comprehensive plant microbiome research. *Front. Ecol. Evol.* 8, 61.
- Salveti, E., Campanaro, S., Campedelli, I., Fracchetti, F., Gobbi, A., Tomielli, G.B., Torriani, S., Felis, G.E., 2016. Whole-metagenome-sequencing-based community profiles of *Vitis vinifera* L. cv. *corvina* berries withered in two post-harvest conditions. *Front. Microbiol.* 7, 937.
- Santos, A., Marquina, D.Y., 2004. 2004. Killer toxin of *Pichia membranifaciens* and its possible use as a biocontrol agent against grey mould disease of grapevine. *Microbiology* 150, 2527–2534.
- Saraf, M., Jha, C.K., Patel, D., 2010. The Role of ACC deaminase producing PGPR in sustainable agriculture. In: Maheshwari, D.K. (Ed.), *Plant Growth and Health Promoting Bacteria. Plant Growth and Health Promoting Bacteria, Microbiology Monographs, Springer Berlin Heidelberg, Berlin, Heidelberg*, pp. 251–272.
- Sarhan, M.S., Patz, S., Hamza, M.A., Youssef, H.H., Mourad, E.F., Fayez, M., Murphy, B., Ruppel, S., Hegazi, N.A., 2018. G3 PhyloChip analysis confirms the promise of plant-based culture media for unlocking the composition and diversity of the maize root microbiome and for recovering unculturable candidate divisions/phyla. *Microbes Environ.* 33 (3), 317–325. [advpub, ME18023](https://doi.org/10.1007/s00203-018-1802-3).
- Sarhan, M.S., Hamza, M.A., Youssef, H.H., Patz, S., Becker, M., ElSawey, H., Nemr, R., Daanaa, H.-S.A., Mourad, E.F., Morsi, A.T., Abdelfadeel, M.R., Abbas, M.T., Fayez, M., Ruppel, S., Hegazi, N.A., 2019. Culturomics of the plant prokaryotic microbiome and the dawn of plant-based culture media—a review. *J. Adv. Res.* 19, 15–27. *Special Issue on Plant Microbiome*.
- Schacherer, J., Shapiro, J.A., Ruderfer, D.M., Kruglyak, L., 2009. Comprehensive polymorphism survey elucidates population structure of *Saccharomyces cerevisiae*. *Nature* 458, 342–345.
- Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Horn, D.J.V., Weber, C.F., 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541.
- Seaver, S.M.D., Liu, F., Zhang, Q., Jeffryes, J., Faria, J.P., Edirisinghe, J.N., Mundy, M., Chia, N., Noor, E., Beber, M.E., Best, A.A., DeJongh, M., Kimbrel, J.A., D’haeseleer, P., McCorkle, S.R., Bolton, J.R., Pearson, E., Canon, S., Wood-Charlson, E.M., Cottingham, R.W., Arkin, A.P., Henry, C.S., 2021. The ModelSEED Biochemistry Database for the integration of metabolic annotations and the reconstruction, comparison and analysis of metabolic models for plants, fungi and microbes. *Nucleic Acids Res.* 49 (D1), D575–D588.
- Seemann, T., 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30, 2068–2069.
- Sereika, M., Kirkegaard, R.H., Karst, S.M., Michaelsen, T.Y., Sørensen, E.A., Wollenberg, R.D., Albertsen, M., 2022. Oxford Nanopore R10.4 long-read sequencing enables the generation of near-finished bacterial genomes from pure cultures and metagenomes without short-read or reference polishing. *Nat. Methods* 19, 823–826.
- Shakir, S., Zaidi, S.S.-A., de Vries, F.T., Mansoor, S., 2021. Plant genetic networks shaping phyllosphere microbial community. *Trends Genet.* 37, 306–316.
- Signorini, M., Borruso, L., Randall, K.C., Dumbrell, A.J., Pii, Y., Mimmo, T., Cesco, S., 2021. Soil heterogeneity within a vineyard impacts the beta but not the alpha microbial agro-diversity. *Appl. Soil Ecol.* 166, 104088.
- Singh, P., Santoni, S., This, P., Péros, J.-P., 2018. Genotype–environment interaction shapes the microbial assemblage in grapevine’s phyllosphere and carposphere: an NGS approach. *Microorganisms* 6, 96.

- Singh, P., Santoni, S., Weber, A., This, P., Péros, J.-P., 2019. Understanding the phyllosphere microbiome assemblage in grape species (Vitaceae) with amplicon sequence data structures. *Sci. Rep.* 9, 14294.
- Sivan, A., Rahimi, O., Lavi, B., Salmon-Divon, M., Weiss, E., Drori, E., Hübner, S., 2021. Genomic evidence supports an independent history of Levantine and Eurasian grapevines. *Plants People Planet* 3, 414–427.
- Smith, N.A., Eamens, A.L., Wang, M.-B., 2011. Viral small interfering RNAs target host genes to mediate disease symptoms in plants. *PLoS Pathog.* 7, e1002022.
- Staudt, G., Kassemeyer, H.H., 1995. Evaluation of downy mildew resistance in various accessions of wild *Vitis* species. *Vitis* 34, 225–228.
- Stefanini, I., Cavalieri, D., 2018. Metagenomic approaches to investigate the contribution of the vineyard environment to the quality of wine fermentation: potentials and difficulties. *Front. Microbiol.* 9, 991.
- Steimetz, E., Trouvelot, S., Gindro, K., Bordier, A., Poinssot, B., Adrian, M., Daire, X., 2012. Influence of leaf age on induced resistance in grapevine against *Plasmopara viticola*. *Physiol. Mol. Plant Pathol.* 79, 89–96.
- Stein, R.R., Bucci, V., Toussaint, N.C., Buffie, C.G., Räscht, G., Pamer, E.G., Sander, C., Xavier, J.B., 2013. Ecological modeling from time-series inference: insight into dynamics and stability of intestinal microbiota. *PLoS Comput. Biol.* 9, e1003388.
- Stempien, E., Pierron, R.J., Adendorff, I., van Jaarsveld, W.J., Halleen, F., Mostert, L., 2020. Host defence activation and root colonization of grapevine rootstocks by the biological control fungus *Trichoderma atroviride*. *Phytopathol. Mediterr.* 59, 615–626.
- Stranska, M., Dzuman, Z., Prusova, N., Behner, A., Kolouchova, I., Lovecka, P., Rezanka, T., Kolarik, M., Hajslova, J., 2022. Fungal endophytes of *Vitis vinifera*—plant growth promoters or potentially toxinogenic agents? *Toxins* 14, 66.
- Sun, X., Kosman, E., Sharon, A., 2020. Stem endophytic mycobiota in wild and domesticated wheat: structural differences and hidden resources for wheat improvement. *J. Fungi* 6, 180.
- Sun, D., Qu, J., Huang, Y., Lu, J., Yin, L., 2021. Analysis of microbial community diversity of muscadine grape skins. *Food Res. Int.* 145, 110417.
- Swift, J.F., Hall, M.E., Harris, Z.N., Kwasniewski, M.T., Miller, A.J., 2021. Grapevine microbiota reflect diversity among compartments and complex interactions within and among root and shoot systems. *Microorganisms* 9, 92.
- Taguiau, J.D., Evallo, E., Balendres, M.A., 2021. *Epicoccum* species: ubiquitous plant pathogens and effective biological control agents. *Eur. J. Plant Pathol.* 159, 713–725.
- Tamaddon-Nezhad, A., Bohan, D., Milani, G.A., Raybould, A., Muggleton, S., 2021. Human-machine scientific discovery. In: *Human-Like Machine Intelligence*. Oxford University Press, pp. 297–315.
- Tedesco, S., Fevreiro, P., Kragler, F., Pina, A., 2022. Plant grafting and graft incompatibility: a review from the grapevine perspective. *Sci. Hortic.* 299, 111019.
- Terral, J.-F., Tabard, E., Bouby, L., Ivorra, S., Pastor, T., Figueiral, I., Picq, S., Chevance, J.-B., Jung, C., Fabre, L., Tardy, C., Compan, M., Bacilieri, R., Lacombe, T., This, P., 2010. Evolution and history of grapevine (*Vitis vinifera*) under domestication: new morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars. *Ann. Bot.* 105, 443–455.
- Theis, K.R., Dheilly, N.M., Klassen, J.L., Brucker, R.M., Baines, J.F., Bosch, T.C., Cryan, J.F., Gilbert, S.F., Goodnight, C.J., Lloyd, E.A., Sapp, J., Vandenkoornhuyse, P., Zilber-Rosenberg, I., Rosenberg, E., Bordenstein, S.R., 2016. Getting the hologenome concept right: an eco-evolutionary framework for hosts and their microbiomes. *Msystems* 1 (2), e00028-16.
- Thibon, C., Shinkaruk, S., Jourdes, M., Bennetau, B., Dubourdieu, D., Tominaga, T., 2010. Aromatic potential of botrytized white wine grapes: Identification and quantification of new cysteine-S-conjugate flavor precursors. *Anal. Chim. Acta* 660, 190–196.

- Thibon, C., Cluzet, S., Mérillon, J.M., Darriet, P., Dubourdieu, D., 2011. 3-sulfanylhexanol precursor biogenesis in grapevine cells: the stimulating effect of *Botrytis cinerea*. *J. Agric. Food Chem.* 59, 1344–1351.
- This, P., Lacombe, T., Thomas, M., 2006. Historical origins and genetic diversity of wine grapes. *Trends Genet.* 22, 511–519.
- Tikhonov, G., Opedal, Ø.H., Abrego, N., Lehtikoinen, A., de Jonge, M.M.J., Oksanen, J., Ovaskainen, O., 2020. Joint species distribution modelling with the r-package Hmsc. *Methods Ecol. Evol.* 11, 442–447.
- Toju, H., Peay, K.G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S., Ushio, M., Nakaoka, S., Onoda, Y., Yoshida, K., Schlaeppli, K., Bai, Y., Sugiura, R., Ichihashi, Y., Minamisawa, K., Kiers, E.T., 2018. Core microbiomes for sustainable agroecosystems. *Nat. Plants* 4, 247–257.
- Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., Singh, B.K., 2020. Plant–microbiome interactions: from community assembly to plant health. *Nat. Rev. Microbiol.* 18, 607–621.
- Trivedi, P., Mattupalli, C., Eversole, K., Leach, J.E., 2021. Enabling sustainable agriculture through understanding and enhancement of microbiomes. *New Phytol.* 230, 2129–2147.
- Trouvelot, S., Bonneau, L., Redecker, D., van Tuinen, D., Adrian, M., Wipf, D., 2015. Arbuscular mycorrhiza symbiosis in viticulture: a review. *Agron. Sustain. Dev.* 35, 1449–1467.
- Tshikantwa, T.S., Ullah, M.W., He, F., Yang, G., 2018. Current trends and potential applications of microbial interactions for human welfare. *Front. Microbiol.* 9, 1156.
- Uyemoto, J.K., Rowhani, A., Ludivisi, D., 2000. An association of rootstock stem lesions in *Vitis* species and different graft-transmissible agents. In: *Proc. 13th ICVG Meet. Adel. Aust.* vol. 83.
- Uyemoto, J.K., Rowhani, A., Luvisi, D., Krag, C.R., 2001. New closterovirus in ‘Redglobe’ grape causes decline of grafted plants. *Calif. Agric.* 55, 28–31.
- Vacher, C., Hampe, A., Porté, A.J., Sauer, U., Compant, S., Morris, C.E., 2016a. The phyllosphere: microbial jungle at the plant–climate interface. *Annu. Rev. Ecol. Evol. Syst.* 47, 1–24.
- Vacher, C., Tamaddoni-Nezhad, A., Kamenova, S., Peyrard, N., Moalic, Y., Sabbadin, R., Schwaller, L., Chiquet, J., Smith, M.A., Vallance, J., Fievet, V., Jakuschkin, B., Bohan, D.A., 2016b. Learning ecological networks from next-generation sequencing data. In: Woodward, G., Bohan, D.A. (Eds.), *Advances in Ecological Research, Ecosystem Services: From Biodiversity to Society, Part 2*. Academic Press, pp. 1–39.
- Vacher, C., Francioni, C., Michel, M., Faivre d’Arcier, J., Fort, T., Chancerel, E., Delmotte, F., Delmas, C.L., (2022), Fungal metabarcoding data for two grapevine varieties (Regent and *Vitis vinifera* L. cv. Cabernet–Sauvignon) inoculated with powdery mildew (*Erysiphe necator*) under drought conditions. *Phytobiomes J.* in press.
- Vandenkoornhuysse, P., Quaiser, A., Duhamel, M., Le Van, A., Dufresne, A., 2015. The importance of the microbiome of the plant holobiont. *New Phytol.* 206 (4), 1196–1206.
- Vass, M., Székely, A.J., Lindström, E.S., Langenheder, S., 2020. Using null models to compare bacterial and microeukaryotic metacommunity assembly under shifting environmental conditions. *Sci. Rep.* 10, 2455.
- Vellend, M., 2010. Conceptual synthesis in community ecology. *Q. Rev. Biol.* 85, 183–206.
- Villano, C., Aversano, R., 2020. Towards grapevine (*Vitis vinifera* L.) mildews resistance: molecular defence mechanisms and new breeding technologies. *Italus Hortus* 27, 1.
- Vitulo, N., Lemos, W.J.F., Calgaro, M., Confalone, M., Felis, G.E., Zapparoli, G., Nardi, T., 2019. Bark and grape microbiome of *Vitis vinifera*: influence of geographic patterns and agronomic management on bacterial diversity. *Front. Microbiol.* 9, 3203.
- Vukicevich, E., Thomas Lowery, D., Úrbez-Torres, J.R., Bowen, P., Hart, M., 2018. Groundcover management changes grapevine root fungal communities and plant–soil feedback. *Plant and Soil* 424, 419–433.



- Waite, H., Whitelaw-Weckert, M., Torley, P., 2015. Grapevine propagation: principles and methods for the production of high-quality grapevine planting material. *N. Z. J. Crop Hortic. Sci.* 43, 144–161.
- Wan, Y., Schwaninger, H.R., Baldo, A.M., Labate, J.A., Zhong, G.-Y., Simon, C.J., 2013. A phylogenetic analysis of the grape genus (*Vitis* L.) reveals broad reticulation and concurrent diversification during neogene and quaternary climate change. *BMC Evol. Biol.* 13, 141.
- Warringer, J., Zörgö, E., Cubillos, F.A., Zia, A., Gjuvsland, A., Simpson, J.T., Forsmark, A., Durbin, R., Omholt, S.W., Louis, E.J., Liti, G., Moses, A., Blomberg, A., 2011. Trait variation in yeast is defined by population history. *PLoS Genet.* 7, e1002111.
- Weiss, S., Van Treuren, W., Lozupone, C., Faust, K., Friedman, J., Deng, Y., Xia, L.C., Xu, Z.Z., Ursell, L., Alm, E.J., Birmingham, A., Cram, J.A., Fuhrman, J.A., Raes, J., Sun, F., Zhou, J., Knight, R., 2016. Correlation detection strategies in microbial data sets vary widely in sensitivity and precision. *ISME J.* 10, 1669–1681.
- Weiss, S., Xu, Z.Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., Lozupone, C., Zaneveld, J.R., Vázquez-Baeza, Y., Birmingham, A., Hyde, E.R., Knight, R., 2017. Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome* 5, 27.
- Wemheuer, F., Taylor, J.A., Daniel, R., Johnston, E., Meinicke, P., Thomas, T., Wemheuer, B., 2020. Tax4Fun2: prediction of habitat-specific functional profiles and functional redundancy based on 16S rRNA gene sequences. *Environ. Microbiome* 15, 11.
- Widder, S., Allen, R.J., Pfeiffer, T., Curtis, T.P., Wiuf, C., Sloan, W.T., Cordero, O.X., Brown, S.P., Momeni, B., Shou, W., Kettle, H., Flint, H.J., Haas, A.F., Laroche, B., Kreft, J.-U., Rainey, P.B., Freilich, S., Schuster, S., Milferstedt, K., van der Meer, J.R., Großkopf, T., Huisman, J., Free, A., Picioreanu, C., Quince, C., Klapper, I., Labarthe, S., Smets, B.F., Wang, H., Soyer, O.S., 2016. Challenges in microbial ecology: building predictive understanding of community function and dynamics. *ISME J.* 10, 2557–2568.
- Wu, M., Zhang, L., Li, G., Jiang, D., Ghabrial, S.A., 2010. Genome characterization of a debilitation-associated mitovirus infecting the phytopathogenic fungus *Botrytis cinerea*. *Virology* 406, 117–126.
- Wu, Y.-W., Simmons, B.A., Singer, S.W., 2016. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32, 605–607.
- Wu, X., Cui, Z., Peng, J., Zhang, F., Liesack, W., 2022a. Genome-resolved metagenomics identifies the particular genetic traits of phosphate-solubilizing bacteria in agricultural soil. *ISME Commun.* 2, 17.
- Wu, X., Rensing, C., Han, D., Xiao, K.-Q., Dai, Y., Tang, Z., Liesack, W., Peng, J., Cui, Z., Zhang, F., 2022b. Genome-resolved metagenomics reveals distinct phosphorus acquisition strategies between soil microbiomes. *mSystems* 7, e01107–e01121.
- Xiao, Y., Angulo, M.T., Friedman, J., Waldor, M.K., Weiss, S.T., Liu, Y.-Y., 2017. Mapping the ecological networks of microbial communities. *Nat. Commun.* 8, 2042.
- Xu, N., Zhang, Z., Shen, Y., Zhang, Q., Liu, Z., Yu, Y., Wang, Y., Lei, C., Ke, M., Qiu, D., Lu, T., Chen, Y., Xiong, J., Qian, H., 2022. Compare the performance of multiple binary classification models in microbial high-throughput sequencing datasets. *Sci. Total Environ.* 837, 155807.
- Yacoub, A., Gerbore, J., Magnin, N., Chambon, P., Dufour, M.-C., Corio-Costet, M.-F., Guyoneaud, R., Rey, P., 2016. Ability of *Pythium oligandrum* strains to protect *Vitis vinifera* L., by inducing plant resistance against *Phaeoconiella chlamydospora*, a pathogen involved in Esca, a grapevine trunk disease. *Biol. Control* 92, 7–16.

- Yergeau, É., Quiza, L., Tremblay, J., 2020. Microbial indicators are better predictors of wheat yield and quality than N fertilization. *FEMS Microbiol. Ecol.* 96, fiz205.
- Yousaf, S., Bulgari, D., Bergna, A., Pancher, M., Quaglino, F., Casati, P., Campisano, A., 2014. Pyrosequencing detects human and animal pathogenic taxa in the grapevine endosphere. *Front. Microbiol.* 5, 327.
- Yu, L., Sang, W., Wu, M.-D., Zhang, J., Yang, L., Zhou, Y.-J., Chen, W.-D., Li, G.-Q., 2015. Novel hypovirulence-associated RNA mycovirus in the plant-pathogenic fungus *Botrytis cinerea*: molecular and biological characterization. *Appl. Environ. Microbiol.* 81, 2299–2310.
- Yue, Y., Huang, H., Qi, Z., Dou, H.-M., Liu, X.-Y., Han, T.-F., Chen, Y., Song, X.-J., Zhang, Y.-H., Tu, J., 2020. Evaluating metagenomics tools for genome binning with real metagenomic datasets and CAMI datasets. *BMC Bioinf.* 21, 334.
- Zaller, J.G., Cantelmo, C., Santos, G.D., Muther, S., Gruber, E., Pallua, P., Mandl, K., Friedrich, B., Hofstetter, I., Schmuckenschlager, B., Faber, F., 2018. Herbicides in vineyards reduce grapevine root mycorrhization and alter soil microorganisms and the nutrient composition in grapevine roots, leaves, xylem sap and grape juice. *Environ. Sci. Pollut. Res.* 25, 23215–23226.
- Zaneveld, J.R., McMinds, R., Vega Thurber, R., 2017. Stress and stability: applying the Anna Karenina principle to animal microbiomes. *Nat. Microbiol.* 2, 1–8.
- Zanne, A.E., Abarenkov, K., Afkhami, M.E., Aguilar-Trigueros, C.A., Bates, S., Bhatnagar, J.M., Busby, P.E., Christian, N., Cornwell, W.K., Crowther, T.W., Flores-Moreno, H., Floudas, D., Gazis, R., Hibbett, D., Kennedy, P., Lindner, D.L., Maynard, D.S., Milo, A.M., Nilsson, R.H., Powell, J., Schildhauer, M., Schilling, J., Treseder, K.K., 2020. Fungal functional ecology: bringing a trait-based approach to plant-associated fungi. *Biol. Rev.* 95, 409–433.
- Zanzotto, A., Morroni, M., 2016. Major biocontrol studies and measures against fungal and oomycete pathogens of grapevine. In: Compant, S., Mathieu, F. (Eds.), *Biocontrol of Major Grapevine Diseases: Leading Research*. CABI, Wallingford, pp. 1–34.
- Zarraonandia, I., Owens, S.M., Weisenhorn, P., West, K., Hampton-Marcell, J., Lax, S., Bokulich, N.A., Mills, D.A., Martin, G., Taghavi, S., van der Lelie, D., Gilbert, J.A., 2015. The soil microbiome influences grapevine-associated microbiota. *MBio* 6, e02527-14.
- Zhang, X., Zhou, Y., Li, Y., Fu, X., Wang, Q., 2017. Screening and characterization of endophytic *Bacillus* for biocontrol of grapevine downy mildew. *Crop Prot.* 96, 173–179.
- Zhang, S., Wang, Y., Chen, X., Cui, B., Bai, Z., Zhuang, G., 2020. Variety features differentiate microbiota in the grape leaves. *Can. J. Microbiol.* 66, 653–663.
- Zhang, J., Liu, Y.-X., Guo, X., Qin, Y., Garrido-Oter, R., Schulze-Lefert, P., Bai, Y., 2021. High-throughput cultivation and identification of bacteria from the plant root microbiota. *Nat. Protoc.* 16, 988–1012.
- Zherdev, A., Vinogradova, S., Byzova, N., Porotikova, E., Kamionskaya, A., Dzantiev, B., 2018. Methods for the diagnosis of grapevine viral infections: a review. *Agriculture* 8, 195.
- Zhou, J., Ning, D., 2017. Stochastic community assembly: does it matter in microbial ecology? *Microbiol. Mol. Biol. Rev.* 81, e00002–e00017.
- Zilber-Rosenberg, I., Rosenberg, E., 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol. Rev.* 32 (5), 723–735.
- Zohary, D., Hopf, M., 1993. Domestication of plants in the old world. In: Zohary, D., Hopf, M. (Eds.), *The Origin and Spread of Cultivated Plants on West Asia, Europe and the Nile Valley*. Clarendon Press, Oxford, England, pp. 143–150.
- Zolti, A., Green, S.J., Sela, N., Hadar, Y., Minz, D., 2020. The microbiome as a biosensor: functional profiles elucidate hidden stress in hosts. *Microbiome* 8, 71.
- Zorrilla, F., Buric, F., Patil, K.R., Zelezniak, A., 2021. metaGEM: reconstruction of genome scale metabolic models directly from metagenomes. *Nucleic Acids Res.* 49, e126.